



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 5 : A61K 48/00, 35/12, 39/00 C12N 15/19, 15/24, 15/25 C12N 15/26, 15/90, 15/63	A1	(11) International Publication Number: WO 93/07906 (43) International Publication Date: 29 April 1993 (29.04.93)
(21) International Application Number: PCT/US92/08999 (22) International Filing Date: 23 October 1992 (23.10.92) (30) Priority data: 781,356 25 October 1991 (25.10.91) US 863,641 3 April 1992 (03.04.92) US (71) Applicant: SAN DIEGO REGIONAL CANCER CENTER [US/US]; 3099 Science Park Road, Suite 200, San Diego, CA 92121 (US). (72) Inventors: SOBOL, Robert, E. ; 5673 La Jolla Hermosa Avenue, La Jolla, CA 92037 (US). FRED, H., Gage ; 8388 Caminito Helecho, La Jolla, CA 92037 (US). ROYSTON, Ivor ; 1515 El Camino del Teatro, La Jolla, CA 92037 (US). FRIEDMAN, Theodore ; 9470 La Jolla Shores Drive, La Jolla, CA 92037 (US). FAKHRAI, Habib ; 1538 Avenida Andante, Oceanside, CA 92056 (US).		(74) Agents: CAMPBELL, Cathryn et al.; Campbell & Flores, 4370 La Jolla Village Drive, Suite 700, San Diego, CA 92122 (US). (81) Designated States: CA, JP, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, SE). Published <i>With international search report.</i>
(54) Title: LYMPHOKINE GENE THERAPY OF CANCER (57) Abstract A novel method of tumor immunotherapy is described comprising the genetic modification of cells resulting in the secretion of cytokine gene products to stimulate a patient's immune response to tumor antigens. In one embodiment, autologous fibroblasts genetically modified to secrete at least one cytokine gene product are utilized to immunize the patient in a formulation with tumor antigens at a site other than an active tumor site. In another embodiment, cells genetically modified to express at least one tumor antigen product and to secrete at least one cytokine gene product are utilized in a formulation to immunize the patient at a site other than an active tumor site.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NL	Netherlands
BE	Belgium	GN	Guinea	NO	Norway
BF	Burkina Faso	GR	Greece	NZ	New Zealand
BG	Bulgaria	HU	Hungary	PL	Poland
BJ	Benin	IE	Ireland	PT	Portugal
BR	Brazil	IT	Italy	RO	Romania
CA	Canada	JP	Japan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	LI	Liechtenstein	SK	Slovak Republic
CI	Côte d'Ivoire	LK	Sri Lanka	SN	Senegal
CM	Cameroon	LU	Luxembourg	SU	Soviet Union
CS	Czechoslovakia	MC	Monaco	TD	Chad
CZ	Czech Republic	MG	Madagascar	TG	Togo
DE	Germany	ML	Mali	UA	Ukraine
DK	Denmark	MN	Mongolia	US	United States of America
ES	Spain			VN	Viet Nam
FI	Finland				

Lymphokine Gene Therapy of CancerBACKGROUND

This application is a continuation-in-part of United States Patent Application Serial No. 07/781,356, filed on October 25, 1991, which is a continuation-in-part of United States Patent Application Serial No. 07/720,872, filed on June 25, 1991, both of which are incorporated herein in their entirety.

Recent advances in our understanding of the biology of the immune system have lead to the identification of important modulators of immune responses, called cytokines (1-3). Immune system modulators produced by lymphocytes are termed lymphokines, a subset of the cytokines. These agents mediate many of the immune responses involved in anti-tumor immunity. Several of these cytokines have been produced by recombinant DNA methodology and evaluated for their anti-tumor effects. The administration of lymphokines and related immunomodulators has resulted in objective tumor responses in patients with various types of neoplasms (4-7). However, current modes of cytokine administration are frequently associated with toxicities that limit the therapeutic value of these agents.

For example, interleukin-2 (IL-2) is an important lymphokine in the generation of anti-tumor immunity (4). In response to tumor antigens, a subset of lymphocytes termed helper T-cells secrete small quantities of IL-2. This IL-2 acts locally at the site of tumor antigen stimulation to activate cytotoxic T-cells and natural killer cells which mediate systemic tumor cell destruction. Intravenous, intralymphatic and intralesional administration of IL-2 has resulted in clinically significant responses in some cancer patients (4-6). However, severe toxicities (hypotension and adema) limit the dose and efficacy of intravenous and intralymphatic IL-

2 administration (5-7). The toxicity of systemically administered lymphokines is not surprising as these agents mediate local cellular interactions and they are normally secreted in only very small quantities.

5 Additionally, other cytokines, such as interleukin-4 (IL-4), alpha interferon (α -INF) and gamma interferon (γ -INF) have been used to stimulate immune responses to tumor cells. Like IL-2, the current modes of administration have adverse side effects.

10 To circumvent the toxicity of systemic cytokine administration, several investigators have examined intralesional injection of IL-2. This approach eliminates the toxicity associated with systemic IL-2 administration (8,9,10). However, multiple intralesional injections are
15 required to optimize therapeutic efficacy (9,10). Hence, these injections are impractical for many patients, particularly when tumor sites are not accessible for injection without potential morbidity.

 An alternative approach, involving cytokine gene
20 transfer into tumor cells, has resulted in significant anti-tumor immune responses in several animal tumor models (11-14). In these studies, the expression of cytokine gene products following cytokine gene transfer into tumor cells has abrogated the tumorigenicity of the cytokine-secreting
25 tumor cells when implanted into syngeneic hosts. The transfer of genes for IL-2 (11,12) γ -INF (13) or interleukin-4 (IL-4) (14) significantly reduced or eliminated the growth of several different histological types of murine tumors. In the studies employing IL-2 gene
30 transfer, the treated animals also developed systemic anti-tumor immunity and were protected against subsequent tumor challenges with the unmodified parental tumor (11,12). Similar inhibition of tumor growth and protective immunity was also demonstrated when immunizations were performed

SUBSTITUTE SHEET

with a mixture of unmodified parental tumor cells and genetically modified tumor cells engineered to express the IL-2 gene. No toxicity associates with localized lymphokine transgene expression was reported in these
5 animal tumor studies (11-14).

While the above gene-transfer procedure has been shown to provide anti-tumor immunity, it still retains practical difficulties. This approach is limited by the inability to transfer functional cytokine genes into many
10 patients' tumor cells, as most patients' tumors cannot be established to grown in vitro and methods for human in vivo gene transfer are not available.

SUMMARY OF THE INVENTION

The present invention demonstrates a novel, more
15 practical method of cytokine cancer immunotherapy. In one approach, selected cells from a patient, such as fibroblasts, obtained, for example, from a routine skin biopsy, are genetically modified to express one or more cytokines. Alternatively, patient cells which may normally
20 serve as antigen presenting cells in the immune system such as macrophages, monocytes, and lymphocytes may also be genetically modified to express one or more cytokines. These modified cells are hereafter called cytokine-expressing cells, ore CE cells. The CE cells are then
25 mixed with the patient's tumor antigens, for example in the form of irradiated tumor cells, or alternatively in the form of purified natural or recombinant tumor antigen, and employed in immunizations, for example subcutaneously, to induce systemic anti-tumor immunity.

30 The cytokines are locally expressed at levels sufficient to induce or augment systemic anti-tumor immune responses via local immunization at sites other than active tumor sites. Systemic toxicity related to cytokine

administration should not occur because the levels of cytokine secreted by the CE cells should not significantly affect systemic cytokine concentrations.

As the amount of cytokine secreted by the CE
5 cells is sufficient to induce anti-tumor immunity but is too low to produce substantial systemic toxicity, this approach provides the benefit of local cytokine administration. In addition, this novel method obviates the need for intralesional injections, which may produce
10 morbidity. Furthermore, the continuous local expression of cytokine(s) at the sites of immunization may also augment anti-tumor immune responses compared to intermittent cytokine injections. This method also provides the advantage of local immunization with the CE cells, as
15 opposed to cumbersome intravenous infusions. This method also eliminates the need for establishing tumor cell lines in vitro as well as transfer of genes into these tumor cells.

This invention also provides an alternative means
20 of localized expression of cytokines to induce and/or increase immune responses to a patient's tumor through genetic modification of cellular expression of both cytokine(s) and tumor antigen(s). In this embodiment, selected cells from a patient are isolated and transduced
25 with cytokine gene(s) as well as gene(s) coding for tumor antigen(s). The transduced cells are called "carrier cells." Carrier cells can include fibroblasts and cells which may normally serve as antigen presenting cells in the immune system such as macrophages, monocytes, and
30 lymphocytes. Transduced carrier cells actively expressing both the cytokine(s) and the tumor antigen(s) are selected and utilized in local immunizations at a site other than active tumor sites to induce anti-tumor immune responses. As with the CE cells, these carrier cells should not
35 produce substantial systemic toxicities, as the levels of

cytokine(s) secreted by the carrier cells should not significantly affect systemic cytokine concentrations. This alternate embodiment is advantageous because it obviates the need to obtain samples of the tumor, which is sometimes difficult. However, carrier cells can be utilized in local immunizations in conjunction with tumor cells, tumor cell homogenates, purified tumor antigens, or recombinant tumor antigens to enhance anti-tumor immunity.

Additionally, this second embodiment retains the same advantages as the first embodiment in that the level of cytokine released by the carrier cells is sufficient to induce anti-tumor immunity but is too low to produce substantial systemic toxicity. In addition, as with the first embodiment, this method obviates the need for intralesional injections, and allows for continuous expression of cytokine(s). This method also eliminates the need for establishing continuous cultures in vitro of tumor cells as well as transfer of genes into these tumor cells, and provides the advantage of local immunization with the carrier cells, as opposed to cumbersome lengthy intravenous infusions.

These approaches may also find application in inducing or augmenting immune responses to other antigens of clinical significance in other areas of medical practice.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows schematic diagrams of retroviral vectors DC/TKIL2, LXSN-IL2, and LNCX-IL2.

Figure 2 shows a mean IL-2 concentration of triplicate supernatant samples measured by ELISA. Supernatants were harvested from overnight cultures of approximately 1.5×10^6 semi-confluent fibroblasts.

SUBSTITUTE SHEET

Figure 3 shows biological activity of the IL-2 secreted by the transduced fibroblasts was demonstrated by measuring mean ^3H -TdR incorporation of an IL-2 dependent T-cell line incubated with triplicate samples of supernatants. Supernatants were harvested from overnight cultures of approximately 1.5×10^6 semi-confluent fibroblasts.

Figure 4 shows comparisons between animals injected with 10^5 CT26 tumor cells alone (\square); 10^5 CT26 tumor cells and 2×10^6 unmodified BALB/C fibroblasts (\blacksquare); 10^5 CT26 tumor cells and 2×10^6 IL-2 transduced BALB/C fibroblasts (\bullet); and 10^5 CT26 tumor cells and 1×10^6 transduced BALB/C fibroblasts (\circ). Tumor measurements are the mean products of the cross-sectional diameter of the tumors from four animals in each treatment group. The (*) indicates statistically significant difference ($P < 0.05$) in tumor growth curves.

Figure 5 shows PCR analysis of neomycin phosphotransferase DNA sequences. Lane 1 - positive control pLXSN-RI-IL2. Lanes 2 through 4 tests genomic DNA; Lanes 5 and 6 ovary genomic DNA; Lane 7 negative control, no DNA. Identical results were obtained with liver, spleen and lung genomic DNA (data not shown).

Figure 6 shows the effect of IL-2 modified fibroblasts on tumor establishment and development using 2×10^6 fibroblasts mixed with 5×10^4 CT26 tumor cells concentrating on the rate of tumor growth.

Figure 7 shows the effect of IL-2 modified fibroblasts on tumor establishment and development using 2×10^6 fibroblasts mixed with 5×10^4 CT26 tumor cells concentrating on the time of tumor onset for the individual animal in each treatment group.

Figure 8 shows the effect of IL-2 modified fibroblasts on tumor establishment and development using 2×10^6 fibroblasts mixed with 1×10^5 CT26 tumor cells concentrating on the rate of tumor growth.

5 Figure 9 shows the effect of IL-2 modified fibroblasts on tumor establishment and development using 2×10^6 fibroblasts mixed with 1×10^5 CT26 tumor cells concentrating on the time of tumor onset for the individual animal in each treatment group.

10 Figure 10 shows the effect of IL-2 modified cells on tumor establishment and development using 2×10^6 DCTK-IL2-modified CT26 tumor cells mixed with 1×10^5 unmodified CT26 compared to 2×10^6 DCTK-IL2-modified fibroblasts mixed with 1×10^5 CT26 concentrating on the rate of tumor growth.

15 Figure 11 shows the effect of IL-2 modified cells on tumor establishment and development using 2×10^6 DCTK-IL2-modified CT26 tumor cells mixed with 1×10^5 unmodified CT26 compared to 2×10^6 DCTK-IL2-modified fibroblasts mixed with 1×10^5 CT26 concentrating on the time of tumor onset
20 for the individual animal in each treatment group.

Figure 12 shows the effect of IL-2 modified fibroblasts on induction of systemic anti-tumor immunity and the rate of tumor growth. Mice were immunized with 2×10^6 fibroblasts mixed with 2.5×10^5 irradiated CT26 tumor
25 cells 7 days prior to challenge with 5×10^4 fresh tumor cells.

Figure 13 shows the effect of IL-2 modified fibroblasts on induction of systemic anti-tumor immunity and the time of tumor onset for the individual animal in
30 each treatment group. Mice were immunized with 2×10^6 fibroblasts mixed with 2.5×10^5 irradiated CT26 tumor cells 7 days prior to challenge with 5×10^4 fresh tumor cells.

Figure 14 shows the effect of IL-2 modified fibroblasts on induction of systemic anti-tumor immunity and the rate of tumor growth. Mice were immunized with 2×10^6 fibroblasts mixed with 2.5×10^5 irradiated CT26 tumor cells 14 days prior to challenge with 5×10^4 fresh tumor cells.

Figure 15 shows the effect of IL-2 modified fibroblasts on induction of systemic anti-tumor immunity and the time of tumor onset for the individual animal in each treatment group. Mice were immunized with 2×10^6 fibroblasts mixed with 2.5×10^5 irradiated CT26 tumor cells 14 days prior to challenge with 5×10^4 fresh tumor cells.

DETAILED DESCRIPTION

A novel method of tumor immunotherapy is described comprising the genetic modification of cells resulting in the secretion of cytokine gene products to stimulate a patient's immune response to tumor antigens. "Gene" is defined herein to be a nucleotide sequence encoding the desired protein. In one embodiment, autologous fibroblasts genetically modified to secrete at least one cytokine gene product are utilized to immunize the patient in a formulation with tumor antigens at a site other than an active tumor site. In another embodiment, cells genetically modified to express at least one tumor antigen gene product and to secrete at least one cytokine gene product are utilized in formulation to immunize the patient at a site other than an active tumor site. Cytokines are preferably expressed in cells which efficiently secrete these proteins into the surrounding milieu. fibroblasts are an example of such cells. Fibroblasts or other cells can be genetically modified to express and secrete one or more cytokines, as described later in this specification.

SUBSTITUTE SHEET

Tumor antigens can be provided by several methods, including, but not limited to the following: 1) CE cells can be transduced with gene(s) coding for tumor antigens. These "carrier cells" are then utilized in patient immunizations. 2) Cloned gene sequences coding for appropriate tumor antigens can be transferred into cells such as fibroblasts or antigen-presenting cells. These cells are then mixed with CE or carrier cells to immunize the patient. 3) Tumor antigens can be cloned in bacteria or other types of cells by recombinant procedures. These antigens are then purified and employed in an immunization with CE and/or carrier cells. 4) Tumor antigens can be purified from tumor cells and used, along with CE or carrier cells, to immunize the patient. 5) Tumor cells may be irradiated or mechanically disrupted and mixed with CE and/or carrier cells for patient immunizations.

This invention encompasses the following steps: (A) isolation of appropriate cells for generation of CE cells or carrier cells; (B) isolation of cytokine genes or isolation of cytokine genes and tumor antigen genes, as well as appropriate marker and/or suicide genes; (C) transfer of the genes from (B) to produce the CE cells or carrier cells; (D) preparation of immunological samples of the patient's tumor antigens or other suitable tumor antigens for immunization with CE or carrier cells; (E) inactivation of the malignant potential of tumor cells if they are used as a source of tumor antigens for immunization; and (F) preparation of samples for immunization. Following are several embodiments contemplated by the inventors. However, it is understood that any means known by those in the art to accomplish these steps will be usable in this invention.

SUBSTITUTE SHEET

(A) Isolation of Cells to Generate CE and Carrier Cells

Cells to be utilized as CE cells and carrier cells can be selected from a variety of locations in the patient's body. For example, skin punch biopsies provide a readily available source of fibroblasts for use in generating CE cells, with a minimal amount of intrusion to the patient. alternatively, these fibroblasts can be obtained from the tumor sample itself. Cells of hematopoietic origin may be obtained by venipuncture, bone marrow aspiration, lymph node biopsies, or from tumor samples. Other appropriate cells for the generation of CE or carrier cells can be isolated by means known in the art. Non-autologous cells similarly selected and processed can also be used.

(B) Isolation of Genes

Numerous cytokine genes have been cloned and are available for use in this protocol. The genes for IL-2, γ -INF and other cytokines are readily available (1-5, 11-14). Cloned genes of the appropriate tumor antigens are isolated according to means known in the art.

Selectable marker genes such as neomycin resistance (Neo^R) are readily available. Incorporation of a selectable marker gene(s) allows for the selection of cells that have successfully received and express the desired genes. Other selectable markers known to those in the art of gene transfer may also be utilized to generate CE cells or carrier cells expressing the desired transgenes.

"Suicide" genes can be incorporated into the CE cells or carrier cells to allow for selective inducible killing after stimulation of the immune response. A gene

such as the herpes simplex virus thymidine kinase gene (TK) can be used to create an inducible destruction of the CE cells or carrier cells. When the CE cells or carrier cells are no longer useful, a drug such as acyclovir or gancyclovir can be administered. Either of these drugs will selectively kill cells expressing TK, thus eliminating the implanted transduced cells. Additionally, a suicide gene may be a gene coding for a non-secreted cytotoxic polypeptide attached to an inducible promoter. When destruction of the CE or carrier cells is desired, the appropriate inducer of the promoter is administered so that the suicide gene is induced to produce cytotoxic polypeptide which subsequently kills the CE or carrier cell. However, destruction of the CE or carrier cells may not be required.

Genes coding for tumor antigen(s) of interest can be cloned by recombinant methods. The coding sequence of an antigen expressed by multiple tumors may be utilized for many individual patients.

20 (C) Transfer of Genes

Numerous methods are available for transferring genes into cultured cells (15). For example, the appropriate genes can be inserted into vectors such as plasmids or retroviruses and transferred into the cells. Electroporation, lipofection and a variety of other methods are known in the field and can be implemented.

One method for gene transfer is a method similar to that employed in previous human gene transfer studies, where tumor infiltrating lymphocytes (TILs) were modified by retroviral gene transduction and administered to cancer patients (16). In this Phase I safety study of retroviral mediated gene transfer, TILs were genetically modified to express the Neomycin resistance (Neo^R) gene. Following

intravenous infusion, polymerase chain reaction analyses consistently found genetically modified cells in the circulation for as long as two months after administration. No infectious retroviruses were identified in these patients and no side effects due to gene transfer were noted in any patients (16). These retroviral vectors have been altered to prevent viral replication by the deletion of viral gag, pol and env genes.

When retroviruses are used for gene transfer, replication competent retroviruses may theoretically develop by recombination between the retroviral vector and viral gene sequences in the packaging cell line utilized to produce the retroviral vector. We will use packaging cell lines in which the production of replication competent virus by recombination has been reduced or eliminated. Hence, all retroviral vector supernatants used to infect patient cells will be screened for replication competent virus by standard assays such as PCR and reverse transcriptase assays (16). Furthermore, exposure to replication competent virus may not be harmful. In studies of subhuman primates injected with a large inoculum of replication competent murine retrovirus, the retrovirus was cleared by the primate immune system (17). No clinical illnesses or sequelae resulting from replication competent virus have been observed three years after exposure. In summary, it is not expected that patients will be exposed to replication competent murine retrovirus and it appears that such exposure may not be deleterious (17).

(D) Preparation of Immunological Samples of the Patient's Tumor Antigens or Purified Recombinant Tumor Antigens

Tumor cells bearing tumor associated antigens are isolated from the patient. These cells can derive either from solid tumors or from leukemic tumors. For solid

tumors, single-cell suspensions can be made by mechanical separation and washing of biopsy tissue (18).

Hematopoietic tumors may be isolated from peripheral blood or bone marrow by standard methods (19).

5 A second variant is the use of homogenates of tumor cells. Such homogenates would contain tumor antigens available for recognition by the patient's immune system upon stimulation by this invention. Either unfractionated cell homogenates, made, for example, by mechanical
10 disruption or by freezing and thawing the cells, or fractions of homogenates preferably with concentrated levels of tumor antigens, can be used.

 Likewise, purified tumor antigens, obtained for example by immunoprecipitation or recombinant DNA methods,
15 could be used. Purified antigens would then be utilized for immunizations together with the CE cells and/or carrier cells described above to induce or enhance the patient's immune response to these antigens.

 In the embodiments employing carrier cells, tumor
20 antigens are available through their expression by the carrier cells. These carrier cells can be injected alone or in conjunction with other tumor antigen preparations or CE cells. Likewise, when CE cells are used, purified recombinant tumor antigen, produced by methods known in the
25 art, can be used.

 If autologous tumor cells are not readily available, heterologous tumor cells, their homogenates, their purified antigens, or carrier cells expressing such antigens could be used.

(E) Inactivation of Tumor Cells

When viable tumor cells are utilized in immunizations as a source of tumor antigens, the tumor cells can be inactivated so that they do not grow in the patient. Inactivation can be accomplished by several methods. the cells can be irradiated prior to immunization (18). This irradiation will be at a level which will prevent their replication. Such viable calls can then present their tumor antigens to the patient's immune system, but cannot multiply to create new tumors.

Alternatively, tumor cells that can be cultured may be transduced with a suicide gene. As described above, a gene such as the herpes simplex thymidine kinase (TK) gene can be transferred into tumor cells to induce their destruction by administration of acyclovir or gancyclovir. After immunization, the TK expressing tumor cells can present their tumor antigens, and are capable of proliferation. After a period of time during which the patients's immune response is stimulated, the cells can be selectively killed. This approach might allow longer viability of the tumor cells utilized for immunizations, which may be advantageous in the induction or augmentation of anti-tumor immunity.

(F) Preparation of Samples for Immunization

CE cells and/or carrier cells and tumor cells, and/or homogenates of tumor cells and/or purified tumor antigen(s), are combined for patient immunization. Approximately 10^7 tumor cells will be required. If homogenates of tumor cells or purified or non-purified fractions of tumor antigens are used, the tumor dose can be adjusted based on the normal number of tumor antigens usually present on 10^7 intact tumor cells. The tumor preparation should be mixed with numbers of CE or carrier

cells sufficient to secrete cytokine levels that induce anti-tumor immunity (11-12) without producing substantial systemic toxicity which would interfere with therapy.

5 The cytokines should be produced by the CE cells or the carrier cells at levels sufficient to induce or augment immune response but low enough to avoid substantial systemic toxicity. This prevents side effects created by previous methods' administration of greater than physiological levels of the cytokines.

10 These mixtures, as well as carrier cells that are utilized alone, will be formulated for injection in any manner known in the art acceptable for immunization. Because it is important that at least the CE cells and carrier cells remain viable, the formulations must be
15 compatible with cell survival. Formulations can be injected subcutaneously, intramuscularly, or in any manner acceptable for immunization.

Contaminants in the preparation which may focus the immune response on undesired antigens should be removed
20 prior to the immunizations.

The following examples are provided for illustration of several embodiments of the invention and should not be interpreted as limiting the scope of the invention.

EXAMPLE IIMMUNIZATION WITH FIBROBLASTS EXPRESSING IL-2
MIXED WITH IRRADIATED TUMOR CELLS5 1) Isolation of Autologous Fibroblasts
for Use in Generating IL-2 Secreting CE Cells

10 Skin punch biopsies will be obtained from each patient under sterile conditions. The biopsy tissue will be minced and placed in RPMI 1640 media containing 10% fetal calf serum (or similar media) to establish growth of the skin fibroblasts in culture. The cultured fibroblasts will be utilized to generate IL-2 secreting CE cells by retroviral mediated IL-2 gene transfer.

2) Retroviral Vector Preparation and
Generation of IL-2 Secreting CE Cells

15 The cultured skin fibroblasts will then be infected with a retroviral vector containing the IL-2 and Neomycin resistance (Neo^R) genes. An N2 vector containing the Neo^R gene will be used, and has been previously utilized by a number of investigators for in vitro and in vivo work,
20 including investigations with human subjects (16). The IL-2 vector will be generated from an N2-derived vector, LLRNL, developed and described by Friedmann and his colleagues (20). It will be made by replacement of the luciferase gene of LLRNL with a full-length cDNA encoding
25 human IL-2. Retroviral vector free of contaminating replication-competent virus is produced by transfection of vector plasmid constructions into the helper-free packaging cell line PA317. Before infection of patients' cells, the vector will have been shown to be free of helper virus. In
30 the event that helper virus is detected, the vector will be produced in the GP + envAM12 packaging cell line in which

the viral gag and pol genes are separated from the env, further reducing the likelihood of helper virus production.

3) Transduction Protocol

The cultured primary fibroblasts will be incubated with supernatant from the packaging cell line as described (20). Supernatant from these cells will be tested for adventitious agents and replication competent virus as described (16) and outlined in Table 1. The fibroblasts are washed and then grown in culture media containing G418, (a neomycin analogue) to select for transduced cells expressing the Neo^r gene. The G418-resistant cells will be tested for expression of the IL-2 gene by measuring the concentration of IL-2 in the culture supernatant by an enzyme linked immunosorbent assay (ELISA) (12). G418-resilient cells expressing IL-2 will be stored at -70°C until required for subsequent use in immunizations.

Table 1

Adventitious Agents and Safety Testing

20	1. Sterility
	2. Mycoplasma
	3. General Safety
	4. Viral Testing
	LCM Virus
25	Thymic agent
	S+/L- eco
	S+/L-xeno
	S+/L- amphi
	3T3 amplification
30	MRC-5/Vero

SUBSTITUTE SHEET

4) Preparation of Irradiated Tumor Cells

Tumors obtained from clinically indicated surgical resections or from superficial lymph node or skin metastases will be minced into 2-3 mm pieces and treated with collagenase and DNase to facilitate separation of the tumor into a single cell suspension. The collected cells will be centrifuged and washed in RPMI 1640 media and then cryopreserved in a solution containing 10% dimethyl sulphoxide and 50% fetal calf serum in RPMI 1640 media. The cells will be stored in liquid nitrogen until the time of administration. Prior to their use in subcutaneous immunizations, the cells will be thawed, washed in media free of immunogenic contaminants, and irradiated with 4,000 rads per minute for a total of 20,000 rads in a cesium irradiator.

5) Patient Selection

Patients will have a histologically confirmed diagnosis of cancer. Patients with tumors that must be resected for therapeutic purposes or with tumors readily accessible for biopsy are most appropriate for this embodiment of the invention.

6) Pretreatment Evaluation

The following pretreatment evaluations will be performed:

- 1) History and physical examination including a description and quantification of disease activity.

2) Performance Status Assessment

- 0 = Normal, no symptoms
1 = Restricted, but ambulatory
2 = Up greater than 50% of waking hours, capable of self-care
3 = Greater than 50% of waking hours confined to bed or chair, limited self-care
4 = Bedridden

3) Pretreatment Laboratory:

CBC with differential, platelet count, PT, PTT, glucose, BUN, creatinine, electrolytes, SGOT, SGPT, LDH, alkaline phosphatase, bilirubin, uric acid, calcium, total protein albumin.

4) Other Analyses:

Urinalysis

CH₅₀, C₃ and C₄ serum complement levels

Immunophenotyping of peripheral blood B cell and T cell subsets

Assays for detectable replication-competent virus in peripheral blood cells

PCR assays of peripheral blood leukocytes for Neo^R, IL-2 and viral env

5) Other Pretreatment Evaluation:

Chest X-ray and other diagnostic studies including computerized tomography (CT), magnetic resonance imaging (MRI) or radionuclide scans may be performed to document and quantify the extent of disease activity.

Follow-up evaluations of these assessments at regular intervals during the course of therapy (approximately every 1 to 3 months) will be useful in determining response to therapy and potential toxicity.

SUBSTITUTE SHEET

permitting adjustments in the number of immunizations administered.

7) Restrictions on Concurrent Therapy

For optimal effects of this treatment, patients
5 should receive no concurrent therapy which is known to suppress the immune system.

8) Final Formulation

Each patient will receive subcutaneous immunizations with a mixture of irradiated tumor cells and autologous fibroblast CE cells genetically modified to secrete IL-2. Approximately 10^7 tumor cells will be mixed with 10^7 fibroblasts known to secrete at least 20 units/ml of IL-2 in tissue culture when semi-confluent (12). The irradiated tumor cells and genetically modified fibroblasts
15 will be placed in a final volume of 0.2 ml normal saline for immunization.

9) Dose Adjustments

At least two subcutaneous immunizations will be administered, two weeks apart, with irradiated tumor cells and autologous fibroblasts genetically modified to secrete IL-2. If no toxicity is observed, subsequent booster immunizations may be administered periodically (at least one week apart) to optimize the anti-tumor immune response.
20

J) Treatment of Potential Toxicity

Toxic side effects are not expected to result from these immunizations. However, potential side effects of these immunizations are treatable in the following manner:
25

SUBSTITUTE SHEET

If massive tumor cell lysis results, any resulting uric acid nephropathy, adult respiratory distress syndrome, disseminated intravascular coagulation or hyperkalemia will be treated using standard methods.

- 5 Local toxicity at the sites of immunization will be treated with either topical steroids and/or surgical excision of the injection site as deemed appropriate.

- Hypersensitivity reactions such as chills, fever and/or rash will be treated symptomatically with antipyretics and antihistamines. Patients should not be treated prophylactically. Should arthralgias, lymphadenopathy or renal dysfunction occur, treatment with corticosteroids and/or antihistamines will be instituted. Anaphylaxis will be treated by standard means such as administration of epinephrine, fluids, and steroids.
- 10
15

EXAMPLE II

A. Retroviral IL-2 Gene Transfer and Expression in Fibroblasts

- Retroviral vectors were employed to transfer and express IL-2 and neomycin phosphotransferase genes in murine and primary human fibroblasts. The retroviral vector DC/TKIL2 produced by Gilboa and co-workers (Gansbacher, et al., J. Exp. Med. 172:1217-1223, 1990, which is incorporated herein by reference) was utilized to transduce murine fibroblasts for application in an animal tumor model (see Section B below). Human fibroblasts were transduced with the retroviral vector LXSN-RI-IL2. Schematic diagrams of the structure of these retroviral vectors are provided in Figure 1. A more complete description of the LXSN-RI-IL2 vector, including its nucleotide sequence, is provided in Example III and in Tables 2, 3 and 4.
- 20
25
30

Following infection with the described vectors and selection for 2-3 weeks in growth media containing the neomycin analogue G418, Balb/c and human embryonic fibroblast culture supernatants were harvested and tested for IL-2 by an enzyme-linked immunosorbent assay (ELISA). Figure 2 depicts the levels of IL-2 secreted by the transduced fibroblasts.

These results can be confirmed using negative control fibroblasts infected with an N2-derived retroviral vector expressing an irrelevant gene such as luciferase or β -galactosidase and studies with adult human fibroblasts.

Biological activity of the IL-2 expressed by the transduced human fibroblasts was confirmed by a cell proliferation bioassay employing an IL-2 dependent T cell line. In this assay, supernatant from the transduced fibroblasts and control unmodified fibroblasts were incubated with the IL-2 dependent T cell line CTIL-2. Incorporation of ^3H -thymidine was measured as an indicator of cell proliferation and IL-2 activity (Figure 3).

B. Efficacy of Transduced Fibroblasts in an Animal Tumor Model

The efficacy of fibroblasts genetically modified to secrete IL-2 was tested in an animal model of colorectal carcinoma. In these studies, the Balb/c CT26 tumor cell line was injected subcutaneously with Balb/c fibroblasts transduced to express IL-2. Control groups included animals injected with 1) a mixture of CT26 tumor cells and unmodified fibroblasts; 2) CT26 tumor cells without fibroblasts and 3) transduced fibroblasts alone. No tumors were detected in 3/8 animals treated with transduced fibroblasts and CT26 cells. In contrast, all untreated control animals (8/8) injected with CT26 tumor cells developed palpable tumors. No tumors were detected in the

animals inoculated with transduced fibroblasts without CT26 tumor cells. The mean CT26 tumor size in Balb/c mice injected with the IL-2 secreting fibroblasts was considerably smaller compared to the control groups (Figure 4). A multivariate non-parametric statistical procedure (Koziol, et al., Biometrics 37:383-390, 1981 and Koziol, et al., Computer Prog. Biomed. 19:69-74, 1984, which is incorporated herein by reference) was utilized to evaluate differences in tumor growth among the treatment groups. The tumor growth curves for the four treatment groups presented in Figure 4 were significantly different ($p=0.048$). Subsequent comparisons between treatment groups revealed a significant difference ($p < 0.05$) in tumor growth between animals injected with CT26 tumor cells alone and animals treated with 2×10^6 transduced fibroblasts and CT26 tumor cells (Figure 4).

EXAMPLE III

A. Project Overview

Lymphokine gene therapy of cancer will be evaluated in cancer patients who have failed conventional therapy. An N2-derived vector containing the neomycin phosphotransferase gene will be used. This vector has been employed by a number of investigators for in vitro and in vivo studies including recently approved investigations with human subjects (Rosenberg et al., N. Eng. J. Med., 323:570-578, 1990). The lymphokine vectors used in this investigation will be generated from the N2-derived vector, LXS_N, developed and described by Miller et al., Mol. Cell Biol. 6:2895, 1986 and Miller et al., BioTechniques 7:980, 1989, which are incorporated herein by reference. The vector LXS_N-RI-IL2 contains human IL-2 cDNA under the control of the retroviral 5' LTR promoter and the neomycin phosphotransferase gene under the control of the SV40 promoter (see Figure 1). The normal human IL-2 leader

sequence has been replaced with a chimeric sequence containing rat insulin and human IL-2 leader sequences (see Tables 2, 3 and 4). This chimeric leader sequence enhances IL-2 gene expression.

- 5 To construct the LXSN-RI-IL2 vector, the bacterial plasmid pBC12/CMV/IL2 (Cullen, B.R., DNA 7:645-650, 1988, which is incorporated herein by reference) containing the full-length IL-2 cDNA and chimeric leader sequence was digested with HindIII and the ends were
- 10 blunted using Klenow polymerase. IL-2 cDNA was subsequently released from the plasmid by digestion with BamHI. The IL-2 fragment was purified by electrophoresis in a 1% agarose gel and the appropriate band was extracted utilizing a glass powder method. Briefly, the gel slice
- 15 was dissolved in 4M NaI at 55°. After cooling to room temperature, 4 μ l of oxidized silica solution (BIO-101, La Jolla, CA) was added to adsorb the DNA. The silica was then washed with a cold solution of 50% ethanol containing 0.1 M NaCl in TE buffer. The DNA was eluted from the
- 20 silica by heating at 55° in distilled H₂O. The purified IL-2 cDNA was then directionally ligated into the HpaI-BamHI cloning sites of the pLXSN vector. A more complete description of the pLXSN-RI-IL2 vector and its partial nucleotide sequence are provided in Tables 2, 3, 4, 5 and
- 25 6.

Table 2

Description of the LXSNI-RI-IL2
from position 1 to 6365

<u>Bases</u>	<u>Description</u>
1-589	Moloney murine sarcoma virus 5' LTR
659-1458	The sequence of the extended packaging signal
1469-2151	IL-2 cDNA with chimeric leader sequence
1469-1718	IL-2 chimeric leader sequence
1647-1718	coding region of the signal peptide
1719-2151	Mature IL-2 coding sequence
2158-2159	Mo mu sarcoma virus end/SV 40 start
2159-2503	Simian virus 40 early promoter
2521-2522	Simian virus DNA end/Tn5 DNA start
2557-3351	Neomycin phosphotransferase
3370-3371	Tn5 DNA end/Moloney murine leukemia virus start
3411-4004	Moloney murine leukemia virus 3' LTR
4073-4074	Moloney murine leukemia DNA end/pBR322 DNA start
4074-6365	Plasmid backbone

Table 3

Enzyme	[# Cuts]	Position(s)
Aat1	[2]	1961, 2481
Aat2	[2]	811, 6295
Acc1	[1]	4252
Acc2	[19]	392, 394, 445, 969, 971, 1193, 2751, 3052, 3084, 3807, 3809, 4081, 4083, 4186, 4527, 5108, 5438, 5931, 6263
Acyl1	[5]	808, 2685, 3860, 5910, 6292
Afl1	[13]	260, 273, 328, 626, 756, 1277, 3201, 3676, 3689, 3744, 4041, 5511, 5733
Afl2	[4]	34, 1064, 1955, 3446
Afl3	[2]	1592, 4480
Aha1	[20]	161, 237, 473, 474, 602, 644, 789, 2689, 2849, 3578, 3653, 3888, 3889, 4017, 4059, 4126, 4161, 4860, 5556, 5907
Aha2	[5]	808, 2685, 3860, 5910, 6292
Aha3	[3]	5239, 5258, 5950
Alu1	[33]	29, 33, 119, 190, 411, 654, 734, 742, 1470, 1486, 1751, 1935, 2003, 2446, 2500, 2791, 3249, 3441, 3445, 3532, 3607, 3826, 4069, 4122, 4141, 4422, 4648, 4738, 4784, 5041, 5562, 5662, 5725
Alw1	[20]	1110, 1414, 1665, 2018, 2147, 2160, 2529, 2553, 2864, 2929, 3110, 4027, 5041, 5127, 5129, 5225, 5226, 5689, 6006, 6010
AlwN1	[4]	231, 3572, 3647, 4896
Aoc1	[2]	847, 1076
Aoc2	[19]	323, 413, 426, 597, 1583, 1721, 2631, 2724, 2798, 2988, 3050, 3739, 3828, 3841, 4012, 4300, 4798, 5959, 6044
Aos1	[2]	2787, 5595
ApaL1	[4]	1717, 4296, 4794, 6040

SUBSTITUTE SHEET

Apy1	[22]	315,	623,	801,	814,	1227,	1252,
		1275,	1295,	1325,	1526,	1536,	1558,
		2196,	2251,	2268,	3072,	3731,	4038,
		4629,	4642				4508,
Aqu1	[6]	241,	472,	1998,	3821,	3854,	3887
Asel	[2]	1801,	5545				
Asp700	[1]	5972					
Asp718	[2]	476,	3891				
AspA1	[1]	1145					
Asul	[29]	169,	200,	245,	260,	273,	328,
		626,	756,	826,	839,	1043,	1254,
		1532,	1649,	3201,	3541,	3586,	3616,
		3676,	3689,	3744,	4041,	5415,	5494,
		5733,	6349				5511,
Ava1	[6]	241,	472,	1998,	3821,	3854,	3887
Ava2	[13]	260,	273,	328,	626,	756,	1277,
		3201,	3676,	3689,	3744,	4041,	5511,
							5733
Ava3	[2]	2232,	2304				
Avr2	[2]	1962,	2482				
Bal1	[3]	658,	1169,	2767			
BamH1	[1]	2152					
Ban1	[9]	318,	476,	1200,	2684,	2719,	3734,
		3859,	3891,	5321			
Ban2	[8]	413,	426,	597,	1583,	3050,	3828,
		3841,	4012				
Bbel	[2]	2688,	3863				
Bbv1	[22]	969,	997,	1738,	2493,	2632,	2758,
		2800,	2816,	2909,	3321,	4060,	4131,
		4372,	4390,	4809,	4899,	4902,	5108,
		5600,	5802				5411,
Bcl1	[1]	2526					
Bgl1	[2]	2435,	5493				
Bsp1286I	[19]	323,	413,	426,	597,	1583,	1721,
		2631,	2724,	2798,	2988,	3050,	3739,
		3841,	4012,	4300,	4798,	5959,	6044

SUBSTITUTE SHEET

28

BspH1	[3]	5200,	6208,	6313		
BspM1	[4]	1501,	2500,	2572,	2953	
BssH2	[4]	392,	443,	3082,	3807	
BstE2	[1]	1145				
BstN1	[22]	315,	623,	801,	814,	1227, 1252,
			1275,	1295,	1325,	1526,	1536, 1558, 1630,
			2196,	2251,	2268,	3072,	3731, 4038, 4508,
			4629,	4642			
BstU1	[19]	392,	394,	445,	969,	971, 1193,
			2751,	3052,	3084,	3807,	3809, 4081, 4083,
			4186,	4527,	5108,	5438,	5931, 6263
BstX1	[1]	2060				
BstY1	[11]	2010,	2152,	2521,	2856,	3102, 5121,
			5132,	5218,	5230,	5998,	6015
Bsu36I	[2]	847,	1076			
Ccrl	[1]	1998				
Cfol	[31]	394,	396,	445,	447,	714, 971,
			2679,	2687,	2751,	2788,	3054, 3084, 3086,
			3314,	3809,	3811,	3862,	4083, 4186, 4216,
			4357,	4390,	4660,	4727,	4827, 5001, 5110,
			5503,	5596,	5933,	6265	
Cfr1	[9]	656,	790,	1167,	1188,	2591, 2765,
			3156,	3183,	5761		
Cfr10I	[3]	3004,	3185,	5453		
Cfr13I	[29]	169,	200,	245,	260,	273, 328,
			626,	756,	826,	839,	1043, 1254, 1277,
			1532,	1649,	3201,	3541,	3586, 3616, 3661,
			3676,	3689,	3744,	4041,	5415, 5494, 5511,
			5733,	6349			
Cvnl	[2]	847,	1076			
Ddel	[23]	75,	165,	191,	282,	553, 847,
			1076,	1348,	1692,	2442,	3348, 3487, 3582,
			3657,	3698,	3879,	3967,	4290, 4755, 5164,
			5330,	5870,	6296		
Dpn1	[30]	95,	1104,	1236,	1421,	1659, 2012,
			2154,	2523,	2528,	2547,	2858, 2936, 3017,
			3026,	3104,	3507,	4021,	5048, 5123, 5134,
			5142,	5220,	5232,	5337,	5678, 5696, 5742,
			6000,	6017,	6053		

SUBSTITUTE SHEET

29

Dra1	[3]	5239, 5258, 5950
Dra2	[4]	328, 1277, 3744, 6349
Eae1	[9]	656, 790, 1167, 1188, 2591, 2765, 3156, 3183, 5761
Eag1	[2]	790, 2591
Eco47I	[13]	260, 273, 328, 626, 756, 1277, 3201, 3676, 3689, 3744, 4041, 5511, 5733
Eco52I	[2]	790, 2591
Eco81I	[2]	847, 1076
EcoN1	[2]	850, 1450
EcoO109I	[4]	328, 1277, 3744, 6349
EcoR1	[1]	1460
EcoR1*	[14]	938, 1037, 1460, 1798, 1805, 1928, 2064, 2121, 2236, 2308, 2400, 5240, 5546, 5801
EcoR2	[22]	313, 621, 799, 812, 1225, 1250, 1273, 1293, 1323, 1524, 1534, 1556, 1628, 2194, 2249, 2266, 3070, 3729, 4036, 4506, 4627, 4640
EcoR5	[4]	137, 213, 3554, 3629
EcoT22I	[2]	2232, 2304
Fdi2	[2]	2787, 5595
Fnu4H1	[41]	793, 967, 983, 986, 1191, 1752, 2430, 2507, 2594, 2646, 2657, 2747, 2752, 2789, 2830, 2917, 2920, 2923, 3159, 3255, 3296, 3310, 4074, 4120, 4217, 4270, 4386, 4404, 4407, 4525, 4680, 4823, 4888, 4891, 5097, 5425, 5614, 5764, 5791, 5886, 6115
FnuD2	[19]	392, 394, 445, 969, 971, 1193, 2751, 3052, 3084, 3807, 3809, 4081, 4083, 4186, 4527, 5108, 5438, 5931, 6263
Fok1	[13]	498, 1198, 1358, 1679, 2333, 2552, 3009, 3034, 3912, 4168, 5339, 5520, 5807
Fsp1	[2]	2787, 5595
Hae2	[4]	2688, 3863, 4358, 4728

SUBSTITUTE SHEET

Hae3	[35]	171,	202,	247,	658,	792,	828,
	840,	1045,	1169,	1190,	1255,	1534,	1650,
	1866,	1961,	2423,	2429,	2438,	2481,	2593,
	2767,	3158,	3185,	3543,	3588,	3618,	3663,
	4495,	4506,	4524,	4958,	5416,	5496,	5763,
	6350						
Hap2	[30]	161,	237,	473,	601,	643,	789,
	2590,	2667,	2689,	2717,	2848,	2938,	3005,
	3186,	3578,	3653,	3888,	4016,	4058,	4126,
	4160,	4687,	4834,	4860,	5050,	5454,	5488,
	5555,	5665,	5907				
Hga1	[8]	455,	707,	960,	1580,	4175,	4591,
	5169,	5899					
HgiA1	[9]	413,	1721,	2798,	2988,	3828,	4300,
	4798,	5959,	6044				
Hha1	[31]	394,	396,	445,	447,	714,	971,
	2679,	2687,	2751,	2788,	3054,	3084,	3086,
	3314,	3809,	3811,	3862,	4083,	4186,	4216,
	4357,	4390,	4660,	4727,	4827,	5001,	5110,
	5503,	5596,	5933,	6265			
HinP1	[31]	392,	394,	443,	445,	712,	969,
	2677,	2685,	2749,	2786,	3052,	3082,	3084,
	3312,	3807,	3809,	3860,	4081,	4184,	4214,
	4355,	4388,	4658,	4725,	4825,	4999,	5108,
	5501,	5594,	5931,	6263			
Hinc2	[1]	5914					
Hind2	[1]	5914					
Hind3	[1]	2498					
Hinf1	[14]	298,	517,	857,	868,	1553,	1814,
	3170,	3304,	3356,	3881,	4380,	4455,	4851,
	5368						
Hpa2	[30]	161,	237,	473,	601,	643,	789,
	2590,	2667,	2689,	2717,	2848,	2938,	3005,
	3186,	3578,	3653,	3888,	4016,	4058,	4126,
	4160,	4687,	4834,	4860,	5050,	5454,	5488,
	5555,	5665,	5907				
Hph1	[11]	1214,	1240,	1817,	2863,	4102,	4111,
	5216,	5443,	5859,	6065,	6100		
Kpn1	[2]	480,	3895				
Mae1	[15]	30,	293,	689,	727,	739,	1452,
	1606,	1893,	1963,	2483,	3442,	3709,	4975,
	5228,	5563					

Mae2	[11]	808,	1139,	1180,	1987,	2801,	2988,
	4233,	5183,	5599,	5972,	6292		
Mae3	[20]	38,	1052,	1080,	1145,	1289,	1478,
	1706,	2805,	3111,	3450,	4134,	4229,	4836,
	4899,	5015,	5298,	5629,	5687,	5840,	6028,
Mbo1	[30]	93,	1102,	1234,	1419,	1657,	2010,
	2152,	2521,	2526,	2545,	2856,	2934,	3015,
	3024,	3102,	3505,	4019,	5046,	5121,	5132,
	5140,	5218,	5230,	5335,	5676,	5694,	5740,
	5998,	6015,	6051				
Mbo2	[17]	444,	1145,	1356,	1575,	1617,	1908,
	1911,	3046,	3256,	3336,	4351,	5142,	5213,
	5968,	6046,	6155,	6351			
Mnl1	[54]	291,	444,	508,	534,	560,	639,
	841,	939,	1227,	1330,	1363,	1369,	1372,
	1378,	1408,	1411,	1426,	1433,	1449,	1559,
	1620,	1909,	1921,	2412,	2418,	2443,	2449,
	2455,	2458,	2470,	2508,	2535,	2599,	2735,
	3092,	3286,	3707,	3859,	3878,	3923,	3948,
	3974,	4054,	4087,	4117,	4379,	4587,	4662,
	4911,	5311,	5392,	5540,	5746,	6339	
Mse1	[22]	35,	1065,	1177,	1207,	1231,	1801,
	1843,	1956,	1971,	2124,	2139,	3447,	4261,
	5186,	5238,	5243,	5257,	5310,	5545,	5584,
	5949,	6321					
Mspl	[30]	161,	237,	473,	601,	643,	789,
	2590,	2667,	2689,	2717,	2848,	2938,	3005,
	3186,	3578,	3653,	3888,	4016,	4058,	4126,
	4160,	4687,	4834,	4860,	5050,	5454,	5488,
	5555,	5665,	5907				
Mst1	[2]	2787,	5595				
Mst2	[2]	847,	1076				
Mval	[22]	315,	623,	801,	814,	1227,	1252,
	1275,	1295,	1325,	1526,	1536,	1558,	1630,
	2196,	2251,	2268,	3072,	3731,	4038,	4508,
	4629,	4642					
Nae1	[1]	3187					
Nar1	[2]	2685,	3860				
Ncil	[20]	161,	237,	473,	474,	602,	644,
	789,	2689,	2849,	3578,	3653,	3888,	3889,
	4017,	4059,	4126,	4161,	4860,	5556,	5907
Ncol	[2]	2389,	3117				

Nde1	[1]	4303						
Nde2	[30]	93, 1102, 1234, 1419, 1657, 2010,						
		2152, 2521, 2526, 2545, 2856, 2934, 3015,						
		3024, 3102, 3505, 4019, 5046, 5121, 5132,						
		5140, 5218, 5230, 5335, 5676, 5694, 5740,						
		5998, 6015, 6051						
Nhe1	[3]	29, 1605, 3441						
Nla3	[26]	61, 1263, 1596, 1649, 1835, 1856,						
		2030, 2230, 2302, 2393, 2559, 2904, 3090,						
		3121, 3147, 3473, 4119, 4224, 4484, 5204,						
		5695, 5705, 5783, 5819, 6212, 6317						
Nla4	[28]	153, 246, 262, 320, 478, 627,						
		758, 827, 959, 1202, 1279, 2154, 2200,						
		2272, 2686, 2721, 3678, 3736, 3861, 3893,						
		4042, 4512, 4551, 5323, 5417, 5458, 5669,						
		6259						
Nsi1	[2]	2232, 2304						
Nsp(7524)1	[8]	1596, 1835, 1856, 2230, 2302, 3090,						
		4119, 4484						
Nsp(7524)2	[19]	323, 413, 426, 597, 1583, 1721,						
		2631, 2724, 2798, 2988, 3050, 3739, 3828,						
		3841, 4012, 4300, 4798, 5959, 6044						
NspB2	[12]	119, 190, 1751, 2158, 2791, 3532,						
		3607, 3989, 4192, 4822, 5067, 6008						
NspH1	[8]	1596, 1835, 1856, 2230, 2302, 3090,						
		4119, 4484						
Paer7I	[1]	1998						
Pal1	[35]	171, 202, 247, 658, 792, 828,						
		840, 1045, 1169, 1190, 1255, 1534, 1650,						
		1866, 1961, 2423, 2429, 2438, 2481, 2593,						
		2767, 3158, 3185, 3543, 3588, 3618, 3663,						
		4495, 4506, 4524, 4958, 5416, 5496, 5763,						
		6350						
Ple1	[7]	865, 1547, 3350, 3889, 4374, 4859,						
		5362						
PpuM1	[3]	328, 1277, 3744						
Pss1	[4]	331, 1280, 3747, 6352						
Pst1	[6]	987, 1163, 1888, 2511, 2738, 5618						
Pvu1	[1]	5743						

Pvu2	[6]	119,	190,	1751,	2791,	3532,	3607
Rsa1	[10]	347,	478,	725,	1342,	1519,	1597,
		2991,	3893,	4288,	5853		
Rsr2	[1]	3201					
Sac1	[2]	413,	3828				
Sau1	[2]	847,	1076				
Sau3A1	[30]	93,	1102,	1234,	1419,	1657,	2010,
		2152,	2521,	2526,	2545,	2856,	2934,
		3024,	3102,	3505,	4019,	5046,	5121,
		5140,	5218,	5230,	5335,	5676,	5694,
		5998,	6015,	6051		5740,	
Sau96I	[29]	169,	200,	245,	260,	273,	328,
		626,	756,	826,	839,	1043,	1254,
		1532,	1649,	3201,	3541,	3586,	3616,
		3676,	3689,	3744,	4041,	5415,	5494,
		5733,	6349			5511,	
Scal	[1]	5853					
ScrF1	[42]	161,	237,	315,	473,	474,	602,
		623,	644,	789,	801,	814,	1227,
		1295,	1325,	1526,	1536,	1558,	1630,
		2251,	2268,	2689,	2849,	3072,	3578,
		3731,	3888,	3889,	4017,	4038,	4059,
		4161,	4508,	4629,	4642,	4860,	5556,
						5907	
Sdu1	[19]	323,	413,	426,	597,	1583,	1721,
		2631,	2724,	2798,	2988,	3050,	3739,
		3841,	4012,	4300,	4798,	5959,	6044
Sec1	[38]	159,	235,	314,	324,	472,	536,
		621,	622,	760,	799,	800,	812,
		1294,	1303,	1323,	1324,	1525,	1557,
		2194,	2266,	2389,	2424,	2433,	2482,
		3117,	3576,	3651,	3730,	3740,	3887,
		4036,	4037,	4640		3950,	
SfaN1	[23]	258,	520,	997,	1657,	2107,	2239,
		2311,	2643,	2898,	2984,	3048,	3114,
		3674,	3934,	4146,	4281,	4317,	4357,
		5629,	5820,	6069		4577,	
Sfil	[1]	2435					
Smal	[2]	474,	3889				
Spel	[1]	726					
Sph1	[4]	1835,	2230,	2302,	3090		

Ssp1	[1]	6177
Sst1	[2]	413, 3828
Stul	[2]	1961, 2481
Styl	[9]	324, 536, 1303, 1962, 2389, 2482, 3117, 3740, 3950
Taq1	[15]	860, 1096, 1407, 1418, 1660, 1999, 2514, 2798, 2954, 2978, 3014, 3176, 3367, 4580, 6024
Thal	[19]	392, 394, 445, 969, 971, 1193, 2751, 3052, 3084, 3807, 3809, 4081, 4083, 4186, 4527, 5108, 5438, 5931, 6263
Tth111I	[6]	465, 877, 1275, 2803, 3880, 4227
Xba1	[2]	1892, 3708
Xho1	[1]	1998
Xho2	[11]	2010, 2152, 2521, 2856, 3102, 5121, 5132, 5218, 5230, 5998, 6015
Xma1	[2]	472, 3887
Xma3	[2]	790, 2591
Xmn1	[1]	5972
Xor2	[1]	5743

Table 4

Enzymes which do not cut LXS NR II.L2:

Acc3	Bgl2	Cla1	Hpa1	Nru1
SnaB1				
Apa1	Bsm1	Dra3	Mlu1	PflM1
Spl1				
Asu2	BspM2	Eco47III	Mro1	Sac2
Sst2				
Ban3	BstB1	Esp1	Not1	Sall

SUBSTITUTE SHEET

Table 5

from 1 to 6365. Numbered from position 1.

L15M11L2	-----1000+-----2000+-----3000+-----4000+-----5000+-----6000+-----
-NusV 5' long ter	----->
(Split)	>
1 to 603 of N11L2	----->
neomycin phosphotransferase	----->
-NusV 3' long ter	----->
signal	----->
Ant1	-----1-----
Ant2	-----1-----
Acc1	-----1-----
Acc2	-----21-----
Acyl1	-----1-----
Afl1	-----1-----
Afl2	-----1-----
Afl3	-----1-----
Aha1	-----1-----
Aha2	-----1-----
Aha3	-----1-----
Alu1	-----1-----
Alu2	-----1-----
Alu3	-----1-----
Acc1	-----1-----
Acc2	-----1-----
Acc3	-----1-----
Apal1	-----1-----
Apal2	-----1-----
Apal3	-----1-----
Apal4	-----1-----
Apal5	-----1-----
Apal6	-----1-----
Apal7	-----1-----
Apal8	-----1-----
Apal9	-----1-----
Apal10	-----1-----
Apal11	-----1-----
Apal12	-----1-----
Apal13	-----1-----
Apal14	-----1-----
Apal15	-----1-----
Apal16	-----1-----
Apal17	-----1-----
Apal18	-----1-----
Apal19	-----1-----
Apal20	-----1-----
Apal21	-----1-----
Apal22	-----1-----
Apal23	-----1-----
Apal24	-----1-----
Apal25	-----1-----
Apal26	-----1-----
Apal27	-----1-----
Apal28	-----1-----
Apal29	-----1-----
Apal30	-----1-----
Apal31	-----1-----
Apal32	-----1-----
Apal33	-----1-----
Apal34	-----1-----
Apal35	-----1-----
Apal36	-----1-----
Apal37	-----1-----
Apal38	-----1-----
Apal39	-----1-----
Apal40	-----1-----
Apal41	-----1-----
Apal42	-----1-----
Apal43	-----1-----
Apal44	-----1-----
Apal45	-----1-----
Apal46	-----1-----
Apal47	-----1-----
Apal48	-----1-----
Apal49	-----1-----
Apal50	-----1-----
Apal51	-----1-----
Apal52	-----1-----
Apal53	-----1-----
Apal54	-----1-----
Apal55	-----1-----
Apal56	-----1-----
Apal57	-----1-----
Apal58	-----1-----
Apal59	-----1-----
Apal60	-----1-----
Apal61	-----1-----
Apal62	-----1-----
Apal63	-----1-----
Apal64	-----1-----
Apal65	-----1-----
Apal66	-----1-----
Apal67	-----1-----
Apal68	-----1-----
Apal69	-----1-----
Apal70	-----1-----
Apal71	-----1-----
Apal72	-----1-----
Apal73	-----1-----
Apal74	-----1-----
Apal75	-----1-----
Apal76	-----1-----
Apal77	-----1-----
Apal78	-----1-----
Apal79	-----1-----
Apal80	-----1-----
Apal81	-----1-----
Apal82	-----1-----
Apal83	-----1-----
Apal84	-----1-----
Apal85	-----1-----
Apal86	-----1-----
Apal87	-----1-----
Apal88	-----1-----
Apal89	-----1-----
Apal90	-----1-----
Apal91	-----1-----
Apal92	-----1-----
Apal93	-----1-----
Apal94	-----1-----
Apal95	-----1-----
Apal96	-----1-----
Apal97	-----1-----
Apal98	-----1-----
Apal99	-----1-----
Apal100	-----1-----

Table 5 (Cont'd)

Table 5 (Cont'd)

from 1 to 6369. Numbered from position 1.

121111112	1000+	2000+	3000+	4000+	5000+	6000+
-NucV 5' long ter						
(Splice)						
1 to 603 of 11112						
macayola phosphatase						
-NucV 3' long ter						
signal	1	2				
Ecot221	111111112	111111112	111111112	111111112	111111112	111111112
Pd12	111111112	111111112	111111112	111111112	111111112	111111112
Pnu4H1	111111112	111111112	111111112	111111112	111111112	111111112
PnuD2	111111112	111111112	111111112	111111112	111111112	111111112
Poh1	111111112	111111112	111111112	111111112	111111112	111111112
Pep1	111111112	111111112	111111112	111111112	111111112	111111112
Mae2	111111112	111111112	111111112	111111112	111111112	111111112
Mae3	111111112	111111112	111111112	111111112	111111112	111111112
Mep3	111111112	111111112	111111112	111111112	111111112	111111112
Mga1	111111112	111111112	111111112	111111112	111111112	111111112
MglA1	111111112	111111112	111111112	111111112	111111112	111111112
Mha1	111111112	111111112	111111112	111111112	111111112	111111112
MlaP1	111111112	111111112	111111112	111111112	111111112	111111112
MlaO2	111111112	111111112	111111112	111111112	111111112	111111112
MlaD2	111111112	111111112	111111112	111111112	111111112	111111112
MlaD3	111111112	111111112	111111112	111111112	111111112	111111112
MlaF1	111111112	111111112	111111112	111111112	111111112	111111112
Mpa2	111111112	111111112	111111112	111111112	111111112	111111112
Mph1	111111112	111111112	111111112	111111112	111111112	111111112
Mpl1	111111112	111111112	111111112	111111112	111111112	111111112
Mae1	111111112	111111112	111111112	111111112	111111112	111111112
Mae2	111111112	111111112	111111112	111111112	111111112	111111112
Mae3	111111112	111111112	111111112	111111112	111111112	111111112
Mbo1	111111112	111111112	111111112	111111112	111111112	111111112
Mbo2	111111112	111111112	111111112	111111112	111111112	111111112
Mu11	111111112	111111112	111111112	111111112	111111112	111111112
Mae1	111111112	111111112	111111112	111111112	111111112	111111112
Mep1	111111112	111111112	111111112	111111112	111111112	111111112

from 1 to 6365. Numbered from position 1.




Table 6 (Cont'd)

360 370 380 390 400 410 420
TTATTGAAC TAACCAATCA GTTGGCTTCT CGCTTCTGTT CGCGGGCTTC CGCTCTCCGA GCTCAATAAA
AATAAACTTG ATTCGTTAGT CAAGCGAAGA GCGAAGACAA GCGCGCGAAG GCGAGAGGCT CGAGTTATTT

>Bam2
>Sac1
>Sst1
>HgiA1
>BssH2

430 440 450 460 470 480 490
AGAGCCCAACA ACCGCTCACT CGCGGGCCCA GTTTCCTCAT AACTGCGCTC GCGCGGGTAC CGGTATTCCG
TCTCGGGTGT TCGCGAGTCA GCGCGGGCT CAGAAGGCTA TCTGACGCAG CGCGCCCATC GGCATAAGCG

>Bam2
>BssH2
>HgiA1
>Tth1111
>Xma1
>Ava1
>Acl
>Sma1
>Rsa1
>Kpn1

500 510 520 530 540 550 560
AATAAAGCT CTTCCTGTTT GCATCCGAAT CGTGTCTCTG CTGTTCTCTG CGAGGGTCTC CTCTGAGTGA
TTATTTCGGA GAAGGACAAA CGTACGCTTA GCAACAGAGC GACAAGGAAC CCTCCAGAG GAGACTCACT

>Sty1

570 580 590 600 610 620 630
TTGACTAACC ACGAGCGGGG TCTTTCATT CGCGGGCTGT CGCGGATTTC GAGAGCCCTG CCGAGCGGAG
AACTGATGGG TGCTGCCCCC AGAAGTAAA CGCGGAGCA GCGGCTAAAC CTCTGGGAC CCGTCCCTGG

>Bam2

>Bcl1

>Cfr1
 .
 >Eae1
 .
 >extended_packaging_signal
 .
 640 650 660 670 680 690 700
 ACCGACCCAC CACCGGAGG TAGCTGGCC AGCAACTTAT CTGTGCTGT COGATTGTCT AGTGTCTATC
 TGGCTGGGTC CTGCGGCTTC ATTGACGGG TCGTTGAATA GACACACACA GGCTAACAGA TCACAGATAC

>Hpa1
 .
 710 720 730 740 750 760 770
 TTTGATGTTA TCGCGCTGCG TCTGTACTAG TTACCTAACT AGCTCTGTAT CTGCGGAGCC CGTGTGCGAA
 AACTACAAT ACGCGGAGCG AGACATGATC AATCGATTGA TCGACACATA GACCGGCTCG GCACCACTT

>Eco521
 .
 >Cfr1
 .
 >Eae1
 .
 >Eag1
 .
 >Eae1
 .
 780 790 800 810 820 830 840
 CTGACGAGTT CTGAACAGCC GCGGCGAACC CTGCGAGAGG TCGACGGGAC TTTGCGGCCC GTTTTGTGCG
 GACTGCTCAA GACTTGTGCG GCGGCGTTCG GACGCTCTCG AGGCTGCGTG AAACCGGCGG CAAAAACAGC

>EcoN1
 .
 >Eau361
 .
 >Aoc1
 .
 >Eau1
 .
 >EcoN11
 .

Table 6 (Cont'd)

```
>Cvn1
  |
>Hst2
  |
850      860      870      880      890      900      910
CCCCACCTGA CCAACGGAGT CGATCTGGAA TCCGACCCCG TCAGGATATG TCGTTCTCGT ACGAGACGAG
GGGCTGGACT CCTTCCTCA GGTACACCTT AGGCTGGGGC AGTCTATAC ACCAAGACCA TCCTCTGCTC

>Hqal
      920      930      940      950      960      970      980
AACCTAAAC AGTTCGGCGC TCGCTCTGAA TTTTCTCTTT CCGTTTGGAA CCGAAGCCGC GCGTCTTCTC
TTGGATTTTG TCAAGGGCGC AGGCAGACTT AAAAAGAAA GCGAAACCTT GCGTTCCGCG CCGAGAACAG

>Pst1
  |
990      1000      1010      1020      1030      1040      1050
TGCTGCAGCA TCGTTCTGTG TTCTCTCTGT CTGACTGTGT TTCTGTATTT GTCTGAAAT TACGGCCAGA
ACGACGTGCT AGCAAGACAC AACAGACACA GACTGACACA AAGACATAA CAGACTTTTA ATCGGGTCT

>Aoc1
>Sau1
>Cvn1
>Hst2
>Bsu361
>Afl2
>Eco811
1060      1070      1080      1090      1100      1110      1120
CTGTTACCAC TCGCTTAAGT TTGACCTTAG GTCACTGGAA AGATGTGGAG CCGATCGCTC ACAAGCAGTC
GACAATGGTG AGCGAATTCA AACTCGAATC CAGTCACTT TCTACAGCTC GCGTACGGAG TTTTGGTCA

>Cfr1
>Aap1
>Eae1
>Eae1
```

Table 6 (Cont'd)

	>Kae2	>BstE2		>Pst1	>Bcl1	>Kae2	>Cfr1
1130	1140	1150	1160	1170	1180	1190	
CGTAGATGTC	AAGAAGAGAC	GTTCGGTTAC	CTTCTGCTCT	CCACAATGCC	CAACCTTTAA	CGTCGGATCG	
CCATCTACAG	TTCTTCTCTG	CAACCCAATG	GAAGACGAGA	CGTCTTACCG	GTTCGAAATT	CCAGCCTAAC	
	>Bam1	>Hph1		>Hph1			
1200	1210	1220	1230	1240	1250	1260	
CGCGGAGACG	CCACCTTTAA	CGGAGACCTC	ATCACCCACG	TTAAGATCAA	GGTCTTTTCA	CCTCGCCCCG	
CGCGCTCTCG	CGTCGAAATT	GGCTCTCGAG	TAGTCGGTCC	AATTCTAGTT	CCAGAAAAGT	CGACCGCCCC	
	>Pss1						
	>Dra2						
	>EcoO1091						
	>PvuK1						
	>Tth1111			>Sty1			
1270	1280	1290	1300	1310	1320	1330	
ATGGACACCC	ACACCAGGTC	CCCTACATCG	TGACCTGCGA	AGCCTTGGCT	TTTGACCCCC	CTCCCTGGCT	
TACCTGTGGG	TCTGGTCCAG	CGGATGTAGC	ACTCGACCTT	TCGGAAACCA	AAACTGGCGG	GAGGGACCCA	
	>Rsa1						
1340	1350	1360	1370	1380	1390	1400	
CAAGCCCTTT	GTACACCTTA	AGCCTCCCGC	TCCTCTTCTT	CCATCCCGCC	CGTCTCTCCC	CGTTCAAGCT	
GTTCGGGAAA	CATGTCCGAT	TCGGAGCGCG	ACGACAACCA	GGTACCGCGC	GCACAGACGG	GGAACTTGCA	
				>EcoM1	>EcoR1		
1410	1420	1430	1440	1450	1460	1470	
CGTGGTTGCA	CGCGGCTCG	ATCTCTCGTT	TATCCAGCCC	TCACTCCTTC	TCTAAGCCCG	AATTGGTTAG	
CGACCAAGCT	CGCGCGGAGC	TAGGAGCGAA	ATAGCTCGCG	ACTGACCAAG	AGATCCCGCC	TTAAGCAATC	

Table 6 (Cont'd)

		>BspM1		>Rsa1			
1480	1490	1500	1510	1520	1530	1540	
CTTCGTAACT	GACCAGCTAC	AGTCGGAAAC	CATCAGCAAG	CAGGTATGTA	CTCTCCAGCG	TCCGCGTGGC	
GAACCAITCA	CTGCTCGATG	TCAGCCTTTC	GTAGTCGTTT	GTCCATACAT	GACAGGTCCC	ACCGCGACCG	
		>Rsa1		>NspM1			
		>Ban2		>Nsp(7524)1			
>P1a1		>Hga1		>Afl3		>Nhe1	
1550	1560	1570	1580	1590	1600	1610	
TTCCCCAGTC	AAGACTCCAG	GCATTTCAGG	CAAGCTCTCG	GCTCTTCTCT	TACATGTACC	TTTTCCTAGC	
AAGGGGTCAG	TTCTGAGGTC	CCTAAACTCC	CTCGGACACC	CGACAAGACA	ATGTACATCG	AAAACGATCG	
1620	1630	1640	1650	1660	1670	1680	
CTCAACCCCTG	ACTATCTTCC	AGGTCAITGT	TCCAACATCG	CCCTGTGGAT	CGACAGGATG	CAACTCTCTGT	
GAGTTCCGAC	TCATAGAAGG	TCCAGTAACA	AGGTTGTACC	GGGACACCTA	GCTGTCTTAC	GTTGAGGACA	
		>HglA1		>Apa1			
1690	1700	1710	1720	1730	1740	1750	
CTTGCAATTGC	ACTAAGTCTT	GCATTGTCTA	CAACAGTCC	ACCTACTTCA	AGTTGTACAA	AGAAAACACA	
GAAAGTAACG	TCATTAGAA	CGTCAACAGT	GTTTGTACCG	TGGATCAAGT	TCAAGATGTT	TCTTTTGTGT	
>Pvu2		>NspB2		>Ase1		>Eph1	
1760	1770	1780	1790	1800	1810	1820	
GCTGCAACTG	GAGCAITTIAC	TGCTCGATT	ACAGATGATT	TTGAATGCAA	TTAATAATTA	CAAGAAATCC	
CGAAGTTTAC	CTGTTAAATG	ACGACCTAAA	TGTCTACTAA	AACITTAOCTT	AATTAATTAAT	GTTCTTAGCG	
>Sph1		>Nsp(7524)1		>NspM1			
>NspM1		>Nsp(7524)1		>Pst1			
1830	1840	1850	1860	1870	1880	1890	
AAACTCAOCC	GCATGCTCAC	ATTAAAGTTT	TACATGCCCC	ACAAGGCCAC	AGAACTGAAA	CATCTGCACT	
TTTGAGTGGG	CGTACGAGTG	TAAATTCAAA	ATGTACCGGT	TCTTCCCGTG	TCTTGACTTT	GTACAGCTCA	

Table 6 (Cont'd)

								>Hsp(7524)1	
								>HspH1	
								>Sph1	
2180	2190	2200	2210	2220	2230	2240			
TGTCAGTTAG	GGTGTGAAA	GTCCCCAGGC	TCCCCAGCAG	GCAGAAGTAT	GCAAGCATC	CATCTCAATT			
ACAGTCAATC	CCACACCTT	CAGGGGTCCG	AGGGGTGCTC	CGTCTTCATA	CGTTTCTAC	GTACAGTTAA			
								>Hcl1	
								>Ava1	
								>EcoT221	
								>Hsp(7524)1	
								>HspH1	
								>Sph1	
2250	2260	2270	2280	2290	2300	2310			
AGTCAGCAAC	CAGGTGTGGA	AAGTCCCEAG	ECTCCCEAGC	AGGCAGAAGT	ATGCAAGCCA	TGCATCTCAA			
TCAGTGGTTG	GTCCACACCT	TTCAGGGGTC	CGAGGGGTCC	TCCGTCTTCA	TACGTTTCT	AGGTAGAGTT			
2320	2330	2340	2350	2360	2370	2380			
TTAGTCAGCA	ACCATAGTCC	CGCCCTTAAC	TCCGCCCATC	CGCCCTCTAA	CTCCGCCCAG	TTCGCCCAT			
AATCAGTGGT	TGGTATCAGG	CGGGCGATTG	AGGGCGGTAG	CGGGCGCATT	GAGGGCGGTC	AAGGGCGGTA			
								>Nco1	
								>Sfi1	
								>Bgl1	
2390	2400	2410	2420	2430	2440	2450			
TCTCGGCGCC	ATGGCTGACT	AATTTTTTIT	ATTATGCCAG	AGGGCGAGCC	CGGCTCGGCG	TCTGAGCTAT			
AGAGGCGGCG	TACGACTGA	TAAAAAATA	TAAATAGCTC	TCCGGCTCCG	CGGAGCGCGG	AGACTGAGTA			
								>Sty1	
								>Ava2	
								>Stu1	
								>BspH1	

55

Table 6 (Cont'd)

2460 2470 2480 2490 2500 2510 2520
 TCCAGAACTA GTCAGGACCG TTTTTCGAG CCTAGGCTT TTGCAAAAAG CTTGGGCTGC ACGTCAGCG
 ACGTCTTCAT CACTCCTCCG AAAAAACCTC CCGATCCGAA AACGTTTTTC GAACCCGACG TCCAGCTCCG

 >BclI
 >Xho2
 >BstYI
 Lmian_virus_DNA_end/Tn5_DNA_start
 2530 2540 2550 2560 2570 2580
 CGATCTGATC AAGAGACACG ATGAGGATCG TTTCGC ATG ATT CAA CAA GAT CGA TTG CAG CCA CGT TGT
 CCTAGACTAG TTCTCTGTCC TACTCCTAGC AAAGCG TAC TAA CTT GTT CTA CCT AAC GTG CGT CCA AGA
 Met Ile Glu Gln Asp Gly Leu His Ala Gly Ser>

>EcoS2I
 >EagI
 >EaeI
 >CfrI
 >Xba3
 2590 2600 2610 2620 2630 2640 2650
 CCG CCG GCT TCG GTG GAG ACG CTA TTC GCG TAT GAC TCG GCA CAA CAG ACA ATC GCG TCG TGT
 GCG CCG CGA ACG CAC CTC TCG GAT AAG CCG ATA CTG ACC CGT GTT GTC TGT TAC CCG ACG ACG
 Pro Ala Ala Trp Val Glu Arg Leu Phe Gly Tyr Asp Trp Ala Glu Glu Thr Ile Gly Cys Ser>

>Hae2
 >BbeI
 >HsrI
 >AcyI
 >Aha2

Table 6 (Cont'd)

>Bsp#1

Table 6 (Cont'd)

2910	2920	2930	2940	2950	2960	
GCT GAT GCA ATG CCG CCG CTC CAT ACG CTT GAT CCG GCT ACC TGC CCA TTC GAG CAC CAA CCG	CGA CTA CGT TAC GCC GCC GAC GTA TGC GAA CTA GCG CGA TGC ACC GGT AAC CTG CTC GTT CCG	Ala Asp Ala Met Arg Arg Leu His Thr Leu Asp Pro Ala Thr Cys Pro Phe Asp His Gln Ala>				
		>RsaI				
		>HgiIA1				
		>Hae2		>Cfr101		
2970	2980	2990	3000	3010	3020	3030
AAA CAT CCG ATC GAG CGA GCA CGT ACT CCG ATG GAA CCG GGT CTT GTC GAT CAG GAT GAT CTC	TTT GTA CCG TAG CTC CCT CGT GCA TGA CCG TAC CTT CCG CCA GAA CAG CTA CTC CTA CTA GAC	Lys His Arg Ile Glu Arg Ala Arg Thr Arg Met Glu Ala Gly Leu Val Asp Gln Asp Asp Leu>				
						>Sph1
						>Hsp(7524)1
		>Bam2			>Bam2	>HepH1
3040	3050	3060	3070	3080	3090	
GAG GAA GAG CAT CAG CCG CTC CCG CCA CCG GAA CTC TTC CCG ACC CTC AAG CCG CCG ATG CCG	CTG CTT CTC GTA CTC CCG GAG CCG GGT CCG CTT GAC AAG CCG TCC GAG TTC CCG CCG TAC CCG	Asp Glu Glu His Gln Gly Leu Ala Pro Ala Glu Leu Phe Ala Arg Leu Lys Ala Arg Met Pro>				
	>Xho2	>Nco1			>Cfr1	
	>BstY1	>Sty1			>Eae1	
3100	3110	3120	3130	3140	3150	
GAG CCG GAG GAT CTC CTC CTC ACC CAT CCG GAT CCG TGC TTG CCG AAT ATC ATG CTC GAA AAT	CTG CCG CTC CTA GAG CAG CAC TCG GTA CCG CTA CCG ACC AAC CCG ATA TAG TAC CAC CTT TTA	Asp Gly Glu Asp Leu Val Val Thr His Gly Asp Ala Cys Leu Pro Asn Ile Met Val Glu Asn>				

Table 6 (Cont'd)

```

      >Cfr101
      |
      >Eae1  >Hae1  >Rsr2
      |      |      |
3160 3170 3180 3190 3200 3210
GGC GGC TTT TCT CGA TTC ATC GAC TGT GCG GCG CTC GGT GTC GCG GAC GCG TAT CAG GAC ATA
CGG GCG AAA AGA CCG AAG TAG CTG ACA CCG GCG GAC CCA CAC GCG CTG GCG ATA CTC CTC TAT
Gly Arg Phe Ser Gly Phe Ile Asp Cys Gly Arg Leu Gly Val Ala Asp Arg Tyr Gln Asp Ile>

3220 3230 3240 3250 3260 3270 3280
GCG TTG GCT ACC CGT GAT ATT GCT GAA GAG CTT GCG GCG GAA TCG GCT GAC GCG TTC CTC GTC
CGC AAC CGA TCG CCA CTA TAA CCA CTT CTC GAA CCG CCG CTT ACC CGA CTG GCG AAG CAG CAC
Ala Leu Ala Thr Arg Asp Ile Ala Glu Glu Leu Gly Gly Glu Trp Ala Asp Arg Phe Leu Val>

3290 3300 3310 3320 3330 3340
CTT TAC GGT ATC GCG GCT CCC GAT TCG CAG GCG ATC GCG TTC TAT GCG CTT CTT GAC CAG TTC
GAA ATC CCA TAG GCG CGA GCG CTA AGC CTC GCG TAG GCG AAG ATA GCG GAA GAA CTC CTC AAG
Leu Tyr Gly Ile Ala Ala Pro Asp Ser Gln Arg Ile Ala Phe Tyr Arg Leu Leu Asp Glu Phe>

>Ple1
|
>Tn5_DNA_end/_Mo-MuLV_DNA_start
3350 3360 3370 3380 3390 3400 3410 3420
TTC TGA GCGGACTC TCGCGTTCG TAAATATAAA CATTTTATTT AGTCTCCACA AAAAGCGCGG AATCAAAGAC
AAG ACT GCGCGTCAG ACCCGAGCT ATTTTATTTT CTAAATATAA TCAGAGCTCT TTTTCCCGCG TACTTTCTG
Phe End>

>Afl2
|
>Nhe1
3430 3440 3450 3460 3470 3480 3490
CCCACTGTA GCTTTGCCAA CCTAGCTTAA GTAAAGCCAT TTTCGAAGGC ATCGAAAAAT ACATAACTCA
GGGTGGACAT CCAAGCGCTT CGATCGAATT CATTCGGGTA AAAGCTTCCG TAACITTTTA TGTATTAECT

>HspB2
|
>Pvu2 >EcoR3
3500 3510 3520 3530 3540 3550 3560
GAATAGAGAA GTTCAGATER AGGTACGGA CAGATCGAAC AGCTGAATAT GCGGCAACA CCATATCTGT
CTTATCTGTT CAAGTCTAGT TCCAGTCTT GTCTACCTTG TCAGCTTATA CCGCGTTTGT CCTATAGACA

```

>ALUN1
 3640 3650 3660 3670 3680 3690 3700
 TCTGTGGTAA GCAGTTCCTG CCCCGGCTCA GGGCCAAGAA CAGATGGTCC CCAGATGCGG TCCAGCGCTC
 AGACACCAAT CGTCAAGGAC GGGGCGGAGT CCGCGTTCTT GTCTACCAGG GGTCTACGGC AGGTCCCGAC

[illegible]

```

                                     >Sac1
                                     .
                                     >Hq1A1
                                     .
                                     >Aqv1
                                     >Stt1
                                     .
                                     >Ava1
                                     >Ban2
                                     .
                                     >BsuH2
                                     .
3780      3790      3800      3810      3820      3830      3840
TGAAGTAAGC AATCAATTGC CTCTCGGCTT CTGTTGCGGC GCTTCTGCTC CCGAGCTCA ATAAAGAGAG
ACTTGATTGC TTAGTCAGGC GAAGAGCGAA GACAAGCGGC CGAAGACGAG GCGCTCGAGT TATTTTCTGC

```

```

>Bae2          >Ban1
      .         ^
>Eba1          >Aap710
      .         |
              >Flc1

```

Table 6 (Cont'd)

		>NotI		>SmaI			
		>BamI		>XmaI		>KpnI	
		>AclI		>AvaI		>RsaI	
>Bam2		>Aha2		>AclI			
3850	3860	3870	3880	3890	3900	3910	
CCACAACCCC	TCACTGGGG	CGCCAGTCCT	CGATTGACT	GAGTGGCCG	GGTACCGTG	TATCCAAATA	
GGTGTGGGG	AGTGAGCCG	CGGCTCAGG	CGCTAACTGA	CTCAGGGGG	CCATCGGCAC	ATAGGTTATT	
		>StyI					
3920	3930	3940	3950	3960	3970	3980	
AACCTCTTC	AGTTCATTC	GACTTGTGG	CTGGCTGTC	CTTCGGAGG	TCTCTCTCA	GTCATTGACT	
TGGGACAAG	TCAAGTAGG	CTGAACACCA	GAGGACAAAG	GAACCTCCG	AGAGGAGACT	CAGTAAGTCA	
		>NotI		>BamI			
4000	4010	4020	4030	4040	4050		
ACCGCTCAG	CGGGCTCTT	CATTTCGGG	CTGCTCCGG	ATCGCCAGG	CGCTGCCCAG	CGACCAACCA	
TGGCACTCG	CCCCACAAA	GTAAACCCC	GAGCAGGCC	TAGCCCTCTG	CGGACCCGTC	CCTGGTGGCT	
		>NotI(7524)1		>NheI		>NheI	
>Mo-MuLV_DNA_end/plasmid_pBR322_DNA_start		>NheI		>NheI		>NheI	
4060	4070	4080	4090	4100	4110	4120	
CCCAACACCG	CGACCTAAG	TGGCTGCTC	CGCGTTTCG	CTGATGACG	TGAAAACCTC	TGACACATTC	
CGGTGCTGG	CTTCATTTC	AACGACCGG	CGCCCAAGC	CAGTACTGC	ACTTTTCCAG	ACTGTGTAGG	
		>HgaI					
4130	4140	4150	4160	4170	4180	4190	
AAGTCCCGCA	GAGGTCACG	GCTTGTCTG	AAGCGGATG	CGGAGGCAG	CAAGCCCGTC	AAGGCGGCTC	
TGAGCGGCT	CTGCACTGT	CGAACAGAC	TTGGCTTAG	CGGCTGTCT	GTTCGGCCAG	TCCCGCCAG	
>NotI		>Tth1111		>KaeI		>AccI	

Table 6 (Cont'd)

4200	4210	4220	4230	4240	4250	4260
AGCGGGTGT	GGCGGGTGT	GGCGGGCAG	CATGACCCAG	TCACGTAGCG	ATAGCGGAGT	GTATACTGGC
TGCGCCACAA	CGCGCCACAG	CGCGCGGTGG	GTACTGGGTC	AGTGCATGGC	TATCGGCTCA	CATATGACCG
4270	4280	4290	4300	4310	4320	4330
TTAACTATCG	GGCATCAGAG	CAGATTGTAC	TCACAGTCCA	CCATATCGGG	TGTGAAATAC	CGCACAGATG
AATTGATAAG	CGTAGTCTC	GTCTAACATG	ACTCTCAGCT	GGTATACGGC	ACACTTTATG	CGTGTCTTAC
4340	4350	4360	4370	4380	4390	4400
CGTAAGGACA	AAATACCGCA	TCAGGCGCTC	TTCGGCTTCC	TGGCTCACTG	ACTCGCTGCG	CTCGGTGCTT
GCATTCTCT	TTTATCGGCT	AGTCCGCGAG	AAGGCGAAGG	ACCGAGTGAC	TGAGCGAAGC	GAGCCACCAA
4410	4420	4430	4440	4450	4460	4470
CGGCTGGCGC	GAGCGGTATC	AGCTCACTCA	AAGCGCGTAA	TACCGTTATC	CACAGAATCA	GGCGATAAGC
GCGGACCGCG	CTCGCCATAG	TGGAGTGAGT	TTCCGGCATT	ATGCCAATAG	GTGTCTTACT	CGGTATTGCG
4480	4490	4500	4510	4520	4530	4540
CAGGAAGCAA	CATGTGACCA	AAAGCGCAGC	AAAGCGCCAG	GAACCGTAAA	AAGGCGCGGT	TGCTGGGCTT
GTCTTTTCTT	GTACACTGCT	TTTCCGGTGG	TTTTCGGGTC	CTTGGCATT	TTCCGGGCGCA	ACGACCGCAA
4550	4560	4570	4580	4590	4600	4610

Table 6 (Cont'd)

TTTCCATAGG CTCGGCCCCC CTCAGGACCA TCACAAAAAT CGACCGTCAA CTCAGCGCTG CCGAAAGCCG
AAAGGTATCC GAGCGCGCGG GACTGCTGCT AGTGTITTTA GCTCGGAGTT CAGTCTCCAC CGCTTTCGGC

4620 4630 4640 4650 4660 4670 4680
ACAGGACTAT AAAGATACCA GCGCTTTCCC CCTCGAAGCT CCGTCTGCG CTCCTCTGTT CCGACCGTGC
TGTCTCATA TTCTATGCT CCGCAAAGCG CGACCTTCCA CGGAGCAGCG CAGAGGACAA CGCTCGGAGC

>Hae2
4690 4700 4710 4720 4730 4740 4750
CGCTTACCG ATACTGTGC GCGTTTCTCC CTTCGGGAG CCGTGGCGCTT TCTCATAGCT CAGCGTTAG
CGGAATGCC TATCGACAG CCGAAGAGG GAAGCGCTTC GCACCGCGAA ACAGTATCGA GTCGGAGTC

>HglAI
>ApaLI
4760 4770 4780 4790 4800 4810 4820
GTATCTCACT TCGGTGAGC TCGTTGCTC CAAGCTGGCG TGTGTCCAG AACCGCGCT TCAGCGCGAC
CATAGAGTCA AGCCACATC AGCAAGCGAG GTTCGACCG ACACACGTC TTGGCGGCA AGTGGCGTC

>HepB2
>P1*1
4830 4840 4850 4860 4870 4880 4890
CGCTGGCGCT TATCGGTAA CTATGCTCT GAGTCCAACC CGGTAAGACA CGACTTATG CCACTCGGAG
GCGACCGCA ATAGCGCAT GATACGAA CTCAGCTTG GCCATTCTGT GCTCAATAG CCGTACCGTC

>AluH1
4900 4910 4920 4930 4940 4950 4960
CAGCCACTCG TAACAGGATT AGCAGACGA CGTATGTAG CCGTCTTACA GAGTTCTGA AGTGGTGGC
GTGGGTGACC ATTCTCTAA TGTCTGCT CCATACATC GCCAGATGT CTCAGACT TCACCACCG

4970 4980 4990 5000 5010 5020 5030
TAACTACCG TAACTAGAA CGACAGTAT TGGTATCTG GCTCTGCTGA AGCAGTTAC CTTCGGAAG
ATTGATCGG ATGTAICTT CCGTCATAA ACCATAGAG CGAGACGACT TGGTCAATG CAAGCGTTT

>HepB2

WO 93/07906

Table 6 (Cont'd)

PCT/US92/08999

5040 5050 5060 5070 5080 5090 5100
 AGAGTTGGTA GCTCTTGATC CCGCAAAACA ACCACCGCTC GTACCGGTGG TTTTITTTGT TCGAAGCAGC
 TCTCAACCAT CGAGAACTAG GCGGTTTGT TCGTGGCGAC CATGCCACC AAAAAACA ACGTTGCTCG

>Xho2 >BstY1
 >BstY1 >Xho2 >Hpa1
 5110 5120 5130 5140 5150 5160 5170
 AGATTACGGC CAGAAAAA GGATCTCAAG AGATCTTTT CATCTTTTCT ACGGGGTCTG ACGCTCAGTG
 TCTAATGGCG GTCTTTTTTT CCTAGAGTTC TTCTAGGAAA CTAGAAAAGA TCGCCACAGC TCGGAGTCAC

>BstY1
 >Xho2 >BstY1 >Dra1
 >Hae2 >BspH1 >Hph1 >Xho2 >Aha3
 5180 5190 5200 5210 5220 5230 5240
 GAACGAAAAC TCACGTTAAG GGATTTTGGT CATCAGATTA TCAAAAAGGA TCTTACCTA GATCTTTTGA
 GTTGTCTTTG AGTCCAATTC CCTAAAACA GTACTCTAAT AGTTTTTCTT ACAAAGTCAT CTAGGAAAAT

>Dra1
 >Aha3
 5250 5260 5270 5280 5290 5300 5310
 AATTAAAAAT GAAGTTTTAA ATCAATCTAA AGTATATATG AGTAAACTTG GTCTGACAGT TACCAATGCT
 TTAATTTTTA CTTCAAAAT TAGTTAGATT TCATATATAC TCATTGGAAC CAGACTGTCA ATGCTTACCA

>Bam1 >Pst1
 5320 5330 5340 5350 5360 5370 5380
 TAATCAGTGA GGCACCTATC TCAGCGATCT GTCTATTTGG TTCAATCATA GTTGGCTGAC TCGGGGTGGT
 ATTAGTCACT CCGTGGATAG AGTGGCTAGA CAGATAAACC AAGTAGGTAT CAACCGACTG ACGGGGAGCA

>BspH1
 5390 5400 5410 5420 5430 5440 5450
 GTAGATAACT ACGATAAGGG AGGGCTTACC ATCTGGGCCC AGTGCTGCAA TGATACGGCG AGACCGACCG
 CATCTATTGA TGCTATGCCC TCCCGAATGG TAGACGGGG TCAAGACGTT ACTATGGCCC TCTGGGTGGG

>Cfr101 >Bgl1
 5460 5470 5480 5490 5500 5510 5520
 TCACCGGCTC CAGATTTATC AGCAATAAAC CAGCCACCGG GAAGGGCCGA CGGCAGAGT CGTCTGCAAA
 AGTGGCGGAG GTCTAAATAG TCGTTATTTG GTGGGTGGCG CTTCGGGGCT CGGTCTTTCA CCAGGAGGTT

Table 6 (Cont'd)

GAATACTCAT ACTCTTCCTT TTTCAATATT ATTGAAGCAT TTATCAGGGT TATTGTCTCA TGAGCCGATA
 CTTATCAGTA TGACAAGCAA AAAGTTATAA TAACTTGGTA AATACTCCCA ATAACAGAGT ACTCGCCTAT

6230 6240 6250 6260 6270 6280 6290

CATATTTGAA TGTATTTAGA AAAATAAACA AATAGCGGTT CCGCCACAT TTCCCCGAAA AGTCCACCT
 GTATAAATT ACATAAATCT TTTTATTGT TTATCCCCAA CCGCGGTGTA AAGCGGCTTT TCAAGGTGGA

>Aat2
 >Aha2
 >Acy1
 >Hae2
 >BspM1
 >EcoO1091
 >Dra2
 6300 6310 6320 6330 6340 6350 6360

CAOGTCTAAG AAACCAATTAT TATCATGACA TTAACCTATA AAAATAGGCG TATCAGGAGG CCGTTTGGTC
 CTGCAGATTC TTTCGTAATA ATAGTACTGT AATTGGATAT TTTTATCCGC ATAGTCTCC GCGAAAGCAG

TTCAA
 AAGTT

types which do not cut L15M1112 :

Ace3	Bgl2	Cla1	Npel	Hru1	SnaB1
Apal	Bam1	Dra3	Nlu1	PflM1	Spl1
Aru2	BspM2	Eco47111	Mro1	Sac2	Set2
Sau3	BstB1	Bsp1	Not1	Sall	

To generate the LXSN-RI-IL2 retroviral vector, 10 micrograms of pLXSN-RI-IL2 DNA was transfected into the ecotropic packaging cell line PE501 by standard calcium phosphate precipitation methods (Miller et al., Mol. Cell Biol. 6:2895, 1986). The transfected PE501 cell line was grown in DMEM medium with 10% FCS. The medium was changed after 24 hours and supernatant harvested 24 hours later to infect the amphotropic packaging cell line PA317 as described (Miller et al., Mol. Cell Biol. 6:2895, 1986 and Miller et al., BioTechniques 7:980, 1989). The infected PA317 cells were harvested by trypsinization 24 hours later and replated 1:20 in DMEM containing 10% FCS and the neomycin analogue G418 (400 μ g/ml). The cells were grown at 37°C in 7% CO₂ atmosphere. The selection medium was changed every 5 days until colonies appeared. On day 14, twenty colonies were selected, expanded and tested for viral production by standard methods (Xu et al., Virology 171:331-341, 1989). Briefly, supernatants were harvested from confluent culture dishes, passed through a .45 μ m filter, diluted with DMEM with 10% FCS and utilized to infect NIH 3T3 cells in the presence of 8 μ g/ml polybrene. After 24 hours, the infected NIH 3T3 cells were grown in culture medium that contained the neomycin analogue G418. After 12-14 days, the colonies were stained, counted and the viral titer calculated as described (Xu et al., Virology 171:331-341, 1989).

Colonies with the highest viral titers ($>10^4$ infectious units/ml) were tested for IL-2 expression by Northern blot analyses. Colonies with the highest viral titers and documented IL-2 expression were cryopreserved and will be utilized as stock cultures to produce the LXSN-RI-IL2 retroviral vector trial.

EXAMPLE IVRETROVIRAL VECTOR CONSTRUCTION AND CYTOKINE EXPRESSION

To increase IL-2 production by transduced cell lines, vectors were used containing different promoters to drive IL-2 expression, and a human IL-2 cDNA was directionally sub-cloned into the insulin secretory signal peptide (17). The IL-2 cDNA was directionally sub-cloned into the parental plasmids of the LXS_N (LTR promoter) and LNCX (CMV promoter) vectors (gifts of Dr. A.D. Miller) (18). The newly constructed vectors (Figure 1), designated as LXS_N-IL2 and LNCX-IL2, were packaged in the PA317 cell line for production of retroviral supernatant. As a control, the high level expressing, double copy vector DC/TKIL-2 vector (thymidine kinase promoter) (a gift of Dr. E. Gilboa) was used for comparison.

These vectors were used to transduce a number of murine and human, primary and established cell lines. Pools of transduced cells were selected and expanded in DMEM medium, containing 10% fetal bovine serum (FBS) and 400 µg/ml of active G-418, a neomycin analogue. The results of expression studies in the MCR9 and Balb/c 3T3 cell lines are presented in Table 7.

Table 7

Comparison of IL-2 expression by fibroblasts
transduced with different IL-2 vectors.

5	Fibroblast	Vector	ng IL-2	Units IL-2
			per 10 ⁶ cells	per day
10	Murine	LNCX (Control)	0.4 ±50%	<1
		LNCX-IL2	33.7 ±11%	67
		LXSN-IL2	6.6 ± 6%	13
		DC/TKIL-2	1.9 ± 5%	4
15	Human	LXSN (Control)	0.7 ±29%	1
		LNCX-IL2	159.5 ±17%	319
		LXSN-IL2	25.5 ±15%	51
		DC/TKIL-2	3.0 ±10%	6

SUBSTITUTE SHEET

EXAMPLE VFIBROBLAST CULTURE AND CONDITIONS FOR RETROVIRAL
TRANSDUCTION

The culture conditions for the growth of primary
5 fibroblasts retroviral transduction were optimized.
Primary fibroblasts were successfully cultured. The
optimal conditions enable the growth of approximately $3-4$
 $\times 10^6$ primary fibroblasts from a 12 mm^2 skin biopsy in
approximately 4-6 weeks. Retroviral infection, G418
10 selection, and expansion of the genetically modified
fibroblasts takes an additional 4-6 weeks.

Exploring the conditions for genetic modification
of primary fibroblasts suggests that optimal transduction
may be obtained by the following procedure: The fibroblasts
15 are synchronized in G1 phase by serum starvation, followed
by stimulation with medium containing 15% fetal bovine
serum 15 hours prior to transduction. The cells are then
subjected to 2 cycles of retrovirus infection, each cycle
lasting approximately 3 hours. The cells are refed with
20 fresh media overnight, and then selection in G418 is
initiated the next day. This method is capable of
transducing 5-15% of the fibroblasts in a culture,
depending on the multiplicity of infection.

This procedure was used to transduce a large
25 number of primary and established fibroblasts. As an
example, Table 8 compares the expression levels of IL-2 in
fibroblast lines transduced with LXS-N-IL2.

SUBSTITUTE SHEET

Table 8

Expression of IL-2 by fibroblasts transduced with LXS^N-IL2.

5	Fibroblast Line	Species	Origin	ng IL-2 Units IL-2	
				per 10 ⁶ cells	per day
	Balb/c 3T3	Murine	Transformed	6.6 ± 6%	13
	MCR9	Human	Embryonic	25.5 ± 15%	51
10	NHDF 313	Human	Skin	25.0 ± 10%	50
	GT1	Human	Skin	15.0 ± 5%	30

These results indicate that the IL-2 expression levels in established, embryonic, and primary fibroblast cultures are similar. Comparison of these data with Table 7 suggest that IL-2 expression is affected more by factors such as different promoters than by the fibroblast line used. Similarly, changes in culture conditions can have important effects on IL-2 expression. Table 9 shows that transduced GT1 cells, a primary human fibroblast culture expressed 15-fold more IL-2 under 100 µg/ml G418 selection than under 25 µg/ml G418 selection. Several other primary fibroblast lines have also been transduced with our vectors and are currently growing under G418 selection.

Table 9

Effect of G418 concentration on IL-2 expression by GT1 cells transduced with LXSN-IL2.

5	Selection dose of G418	ng IL-2 secreted per 10 ⁶ cells per day
10	25 µg/ml	1.0 ± 10%
	50 µg/ml	3.0 ± 6%
	100 µg/ml	15.0 ± 5%

*After three weeks of G418 selection.

EXAMPLE VI

15 COMPARISON OF IL-2 EXPRESSION LEVELS INDUCED PERIPHERAL BLOOD LYMPHOCYTES AND GENETICALLY MODIFIED FIBROBLASTS

In order to compare the production of IL-2 by genetically modified fibroblasts to that achieved by stimulating normal human peripheral blood lymphocytes (nPBL) in vitro, nPBL were isolated by Ficoll-Paque density centrifugation, and cultured in the presence of allogeneic nPBL (mixed lymphocyte culture, MLC) or 2 µM calcium ionophore (CI) (A23187) free acid) plus 17 nM phorbol 12-myristate 13-acetate (PMA). The results of this experiment, present in Table 10, indicate that the level of IL-2 expression in the PMA/CI stimulated normal T cell population was 2 ng/10⁶ cells/24 hours. This is equivalent to IL-2 expression by Balb/c 3T3 fibroblasts transduced with DC/TKIL-2 (Table 7), our least productive vector. The level of IL-2 expression in the MLC was 130 pg/10⁶ cells/24 hours. This was lower than the PMA/CI stimulated culture, presumably because PMA/CI induced a nonspecific response

SUBSTITUTE SHEET

while MLC resulted in specific Th stimulation. When the estimated percentage of antigen-specific Th in the MLC-stimulated population is taken into consideration, the level of IL-2 expression per stimulated T cell becomes
5 equivalent for both methods.

Table 10
Levels of IL-2 secretion by different cells.

	Cells	pg IL-2 secreted per 10 ⁶ cells per day		
10				
	Lymphocytes:			
	Control (non-activated)	5	±	50%
	PMA + Calcium Ionophore	2,000	±	6%
15	Mixed lymphocyte culture	130	±	90%
	Transduced fibroblasts:			
	MCR9-LXSN-IL2	24,000	±	5%
	MCR9-LNCX-IL2	162,000	±	20%
	MCR9-DC/TKIL-2	10,000	±	6%
20				

EXAMPLE VII

FIBROBLAST MEDIATED CYTOKINE GENE THERAPY IN MURINE TUMOR MODELS

Two experimental protocols were used to study the
25 efficacy of fibroblast-mediated cytokine gene therapy on
induction of anti-tumor immunity. The first protocol was
designed to test the effects of genetically modified
fibroblasts on tumor implantation, while the second
protocol was designed to induce a systemic anti-tumor
30 immunity. The results of each experiment are presented
with two figures and one table. In the first figure, the
rate of tumor growth for each treatment group is presented

SUBSTITUTE SHEET

as the mean tumor size in the group over time. In the second figure, a Kaplan-Meier curve presents the time of tumor onset for the individual animals in each treatment group. The number of animals, the number and percentage of tumor free animals, and the tumor size distribution patterns for each experiment are presented in a table.

EXAMPLE VII(a)

EFFECT OF FIBROBLAST MEDIATED CYTOKINE GENE
THERAPY ON TUMOR IMPLANTATION

10 Mice were injected subcutaneously with mixtures
of 5×10^4 CT26 cells and 2×10^6 fibroblasts genetically
modified by different retroviral vectors to express IL-2.
In the control arms injected with tumor cells only, or with
tumor cells mixed with unmodified fibroblasts, 31 of 33
15 animals (94%) developed tumors by 4 weeks (Figures 6 and 7,
Table 9). In contrast, 22 out of the 34 animals (65%)
receiving fibroblast mediated cytokine gene therapy were
tumor free at 3 weeks, and 5 animals (18%) remain tumor
free after 12 weeks. Those animals that received
20 fibroblast mediated IL-2 therapy and developed tumor were
characterized by a delayed onset and rate of tumor growth.

Table 11

Effect of IL-2 modified fibroblasts on tumor establishment and development.
 2 X 10⁶ fibroblasts mixed with 5 X 10⁴ CT26 tumor cells at time of injection.

Fibroblasts mixed with tumor cells	Animal Number		Percent Tumor-free	Tumor Size (mm ²)				Median Tumor Size (mm ²)
	Total	Tumor- free		25-100	101-200	201-300	>301	
After 12 Weeks:*								
Control (no fibroblasts)	11	0	11	0%	1	0	1	420 ± 145
Unmodified fibroblasts**	13	2	11	15%	1	0	1	388 ± 265
DCTK-IL2 fibroblasts	13	0	13	0%	1	3	5	267 ± 168
LNCX-IL2 fibroblasts	13	5	8	39%	5	2	0	72 ± 90

* Mean tumor size is for 4 weeks, the last timepoint at which tumors were measured.

** Two mice in this arm developed intraperitoneal tumors which were not measurable.

After 3 weeks the mean tumor size (measured as the product of the longest and widest tumor axes) in the control group of mice was 128 mm², compared to 68 and 7 mm² in groups of mice injected with tumor cells mixed with fibroblasts transduced with DC/TKIL-2 or LNCX-IL2, respectively. This resulted in a highly significant difference (corrected $\chi^2 = 18.69$, $p = 0.001$) between the IL-2 treated animals compared to the mice treated with CT26 alone or CT26 mixed with unmodified fibroblasts. After four weeks the equivalent measurements were 373, 300 and 72 mm² (Table 11). It is notable that LNCX-IL2, the highest expressing vector caused substantially greater inhibition of tumorigenicity than the lower expressing vector DC/TKIL-2. A multivariate non-parametric statistical procedure (19,20), utilized to evaluate differences in tumor growth, demonstrated that after 4 weeks the differences between the growth curves for the four groups presented in Figure 2 were highly significant ($p < 0.001$). Subsequent comparisons between the control arm and animals that received tumor cells mixed with IL-2 transduced fibroblasts revealed a significant difference ($P < 0.05$). The differences between the animals injected with tumor cells alone, and those injected with tumor cells plus unmodified fibroblasts were not significant, while the differences between animals receiving low IL-2 expressing fibroblast, and those receiving high IL-2 expressing fibroblasts was significant ($P = 0.05$).

When mice were injected with 2×10^6 modified fibroblasts mixed with 1×10^5 live tumor cells the results became more striking (see Figures 8 and 9, and Table 12). All the control animals developed tumors after 4 weeks whereas 33% and 27% of the animals treated with fibroblasts modified with the DCTK-IL2 or LKSN-IL2 vectors (respectively) remain tumor free after 7 weeks (the experiment is ongoing). More dramatically, 75% of the animals treated with fibroblasts modified with the highest

IL-2 producing vector, LNCX-IL2, remain tumor free after 7 weeks. These data clearly demonstrate the importance of an initial high dose of IL-2 to prevent tumor establishment.

SUBSTITUTE SHEET

Table 12

Effect of IL-2 modified fibroblasts on tumor establishment and development.
2 X 10⁵ fibroblasts mixed with 1 X 10⁵ CT26 tumor cells at time of injection.

Fibroblasts mixed with tumor cells	Animal Number		Percent Tumor-free	Tumor Size (mm ²)				Mean Tumor Size (mm ²)
	Total	Tumor- free		25-100	101-200	201-300	> 301	
After 6 Weeks:								
Control (no fibroblasts)**	13	0	13	0	5	2	5	315 ± 197
Unmodified fibroblasts**	20	0	20	0	2	3	14	350 ± 100
DCTK-IL2 fibroblasts	12	4	8	0	1	4	3	185 ± 141
LXSN-IL2 fibroblasts***	15	4	11	0	5	1	2	135 ± 121
LNCX-IL2 fibroblasts	8	6	2	2	0	0	0	8 ± 14

* Mean tumor size is for 4 weeks, the last timepoint at which tumors were measured.

** One mouse in each of these arms developed an intraperitoneal tumor which was not measurable.

*** Three mice in this arm developed intraperitoneal tumors which were not measurable.

- As an additional control, mice were injected with CT26 cells genetically modified to express IL-2 (results not shown). Injection of up to 1×10^6 IL-2 expressing tumor cells into Balb/c mice failed to produce tumors.
- 5 Injection of higher numbers however, resulted in some animals developing tumors with delayed onset. These data confirm the results reported in the literature (1). In order to compare the efficacy of IL-2 producing fibroblasts to IL-2 producing tumor cells, we mixed 2×10^6 CT26 tumor
- 10 cells modified with the DCTK-IL2 vector with 1×10^5 unmodified tumor cells. Figures 10 and 11, and Table 13 show that DCTK-IL2 modified tumor cells are somewhat effective in preventing tumor development. Four weeks after injection, the mean tumor size for the treatment arm
- 15 is 303 mm^2 , compared to 620 mm^2 for the control arm. After 22 weeks, one animal (10%) remains tumor free, compared to none in the control arms. Data for animals treated under the same conditions with DCTK-IL2 modified fibroblasts in a separate experiment are included for comparison purposes.
- 20 This comparison suggests that DCTK-IL2 modified tumor cells have an effect on tumor establishment similar to that of DCTK-IL2 modified fibroblasts.

Table 13

Effect of IL-2 modified cells on tumor establishment and development.
 2 X 10⁶ DCTK-IL2-modified CT26 tumor cells mixed with 1 X 10⁵ CT26 cells compared to 2 X 10⁶ DCTK-IL2-modified fibroblasts mixed with 1 X 10⁵ CT26.

Cells mixed with tumor cells	Animal Number		Percent Tumor-free	Tumor Size (mm ²)				Mean Tumor Size (mm ²)
	Total	Tumor-free		25-100	101-200	201-300	> 301	
After 22 Weeks:*								
Control (no fibroblasts)	5	0	5	0	0	0	5	620 ± 190
Unmodified fibroblasts	5	0	5	0	0	0	5	587 ± 69
DCTK-IL2-modified CT26 cells	10	1	9	1	0	2	5	303 ± 179
DCTK-IL2-modified fibroblasts	8	2	6	0	1	2	3	214 ± 158

* Mean tumor size is for 4 weeks, the last timepoint at which tumors were measured.

EXAMPLE VII(b)EFFECT OF FIBROBLAST MEDIATE CYTOKINE GENE THERAPY
ON SYSTEMIC ANTI-TUMOR IMMUNITY

Groups of Balb/c mice were immunized with
5 2.5 x 10⁵ irradiated tumor cells either alone or mixed with
2 x 10⁶ transduced or unmodified fibroblasts, and challenged
one week later with 5 x 10⁴ live tumor cells in the opposite
flank. These results (Figures 12 and 13, and Table 14)
demonstrate that immunization with irradiated tumor cells
10 and transduced fibroblasts protect some animals against a
live tumor challenge, but that the protection is only
slightly better than that achieved by immunization with
irradiated tumor cells alone or irradiated tumor cells
mixed with unmodified fibroblasts.

SUBSTITUTE SHEET

Table 14

Effect of IL-2 modified fibroblasts on induction of systemic anti-tumor immunity.
Mice immunized with 2×10^6 fibroblasts mixed with 2.5×10^5 irradiated CT26 tumor cells 7 days prior to challenge with 5×10^4 fresh tumor cells.

Fibroblasts mixed with irradiated tumor cells	Animal Number		Percent Tumor-free		Tumor Size (mm ²)			Mean Tumor Size (mm ²)
	Total	Tumor-free	Tumor-bearing		25-100	101-200	201-300 > 301	
After 22 Weeks:								
Control (saline)	20	0	20	0%	0	0	1	19
Irradiated CT26 only**	16	5	11	31%	2	1	2	5
Irradiated CT26 mixed with unmodified fibroblasts	15	4	11	27%	0	1	3	7
DCTK-IL2 fibroblasts**	25	10	15	40%	4	1	1	8
								574 ± 160
								250 ± 277
								266 ± 199
								172 ± 194

* Mean tumor size is for 4 weeks, the last timepoint at which tumors were measured.

** One mouse in each of these arms developed an intraperitoneal tumor which was not measurable.

In a second protocol similar to the one described above, animals were challenged with fresh tumor cells two weeks following immunization with irradiated tumor cells mixed with fibroblasts. The results, shown in Figures 14 and 15, and in Table 15, demonstrate that DCTK-IL2 modified fibroblasts mixed with irradiated tumor cells confers superior protection to subsequent tumor challenge than irradiated tumor cells alone, irradiated tumor cells mixed with unmodified fibroblasts, or irradiated tumor cells mixed with LNCX-modified fibroblasts. After 7 weeks, seven of ten animals (70%) treated with DCTK-IL2 modified fibroblasts remain tumor free compared to only one third of the control animals. At four weeks, the mean tumor size of this group was 41 mm², compared to 180, 170, and 140 mm² for the three control groups. Animals treated with LNCX-IL2 modified fibroblasts were also protected against subsequent tumor challenge, but the results were less striking. In this group, 54% of the animals remain tumor free and the mean tumor size for the group at four weeks was 86 mm². The number of tumor free animals in the group treated with LXSN-IL2 modified fibroblasts was similar to the control groups, although the tumors were slightly delayed in their onset. A multivariate non-parametric statistical procedure (19, 20), utilized to evaluate differences in tumor onset, demonstrated that the differences for the six arms presented in Figure 15 were significant ($p = 0.012$). It further showed that the saline control arm and the arms that received irradiated tumor cells alone or mixed with unmodified or LNCX vector modified fibroblasts formed a statistical group. A second, distinct statistical group was formed by the three arms that received IL-2 vector modified fibroblasts mixed with irradiated tumor cells. Subsequent comparisons between the saline injected control arm and animals that received tumor cells mixed with IL2 transduced fibroblasts revealed a significant difference for all vectors ($p < 0.05$).

Table 15

Effect of IL-2 modified fibroblasts on induction of systemic anti-tumor immunity.
Mice immunized with 2×10^6 fibroblasts mixed with 2.5×10^5 irradiated CT26 tumor cells 14 days prior to challenge with 5×10^4 fresh tumor cells.

Immunization by fibroblasts mixed with irradiated tumor cells	Animal Number			Percent Tumor-free	Tumor Size (mm ²)				Mean Tumor Size (mm ²)
	Total	Tumor- free	Tumor- bearing		25-100	101-200	201-300	>301	
After 7 Weeks:									
Control (saline)**	8	1	7	13%	0	2	1	3	245 ± 173
Irradiated CT26 only	10	3	7	30%	0	2	4	1	180 ± 155
Irradiated CT26 mixed with unmodified fibroblasts	6	2	4	33%	0	2	1	1	170 ± 160
Irradiated CT26 mixed with LNCX-modified fibroblasts	10	3	7	30%	3	0	1	3	140 ± 142
Irradiated CT26 mixed with LNCX-IL2-modified fibroblasts	13	7	6	54%	1	3	1	1	86 ± 112
Irradiated CT26 mixed with LXSN-IL2-modified fibroblasts	12	4	8	33%	5	0	2	1	111 ± 145
Irradiated CT26 mixed with DCTK-IL2-modified fibroblasts	10	7	3	70%	1	2	0	0	41 ± 75

* Mean tumor size is for 4 weeks, the last timepoint at which tumors were measured.

** One mouse in this arm developed an intraperitoneal tumor which was not measurable.

These results demonstrate the feasibility of using genetically modified fibroblasts as a means of delivering cytokine gene therapy. In all experiments, the LNCX-L2 vector proved superior in preventing tumor establishment while the DCTK-IL2 vector was better in the induction of systemic protection against subsequent tumor challenges. These contrasting effects, although somewhat surprising, can be explained by the observation that the CMV promoter is turned off in vivo five days after implantation while the TK promoter remains active for a longer period of time. The implication of this finding is that to apply this method of gene therapy successfully we have to use promoters that result in high level, sustained expression of IL-2 in vivo in the transduced fibroblasts.

The data obtained from this research effort has important implications for all cytokines that have either direct or indirect anti-tumor effects. Furthermore, this data suggests that anti-tumor efficacy is IL-2 dose dependent. Hence, construction of vectors which result in higher levels of cytokine secretion will be a significant advance toward the application of this method of gene therapy.

Reference numbers in parenthesis in the above examples correspond to the following list of references and are incorporated herein by reference.

References

1. Gabrilove, J.L. et al., Monogr. J. Natl. Cancer Inst. 10:73-7 (1990).
- 5 2. Kelso, A., Current Opinion in Immunology, 2:215-25 (1989).
3. Borden, E.C. et al., Cancer, 65:800-14 (1990).
4. Rosenberg, S.A. et al., Ann. Intern. Med., 108:853-864 (1988).
- 10 5. Lotze, M.T. et al., JAMA, 256:3117-3124 (1986).
6. Pizza, G. et al., Lymphokine Research, 7:45-8 (1988).
7. Sarna, G. et al., Journal of Biological Response Modifiers, 9:81-6 (1990).
- 15 8. Gandolfi, L. et al., Hepato-Gastroenterology, 36:352-6 (1989).
9. Bubenik, J. et al., Immunol. Letters, 19:279-82 (1988).
10. Bubenik et al., Immunol. Letters, 23:287-292 (1990).
- 20
11. Fearon, E.R. et al., Cell, 60:387-403 (1990).
12. Gansbacher, B. et al., J. Exp. Med., 172:1217-1224 (1990).
13. Watanabe, Y. et al., Proc. Natl. Acad. Sci., 86:9456-9460 (1989).
- 25

SUBSTITUTE SHEET

14. Tepper, R.I. et al., Cell, 57:503-512 (1989).
15. Kriegler, M., Gene Transfer and Expression: A Laboratory Manual, Stockton Press (1990).
16. Rosenberg, S.A. et al., N. Eng. J. Med., 370
5 (1990).
17. Cornetta, K. et al., Prog. Nucl. Acid Res. Mol. Biol., 36:311-22 (1989).
18. Hoover, H.C. et al., Cancer Res., 44:1671-76 (1984).
- 10 19. Sobol et al. New Eng. J. Med. 316:1111-1117 (1987).
20. Li Xu, et al., Virology, 171:331-341 (1989).

Although the invention has been described with reference to the presently-preferred embodiment, it should be understood that various modifications can be made without departing from the spirit of the invention.

5 Accordingly, the invention is limited only by the following claims.

WE CLAIM:

1. A method of treating cancer in a patient comprising the stimulation of that patient's immune response against the cancer by immunizing said patient at a site other than an active tumor site with a formulation comprising tumor antigens and CE cells genetically modified to express at least one cytokine gene product.
2. The method of claim 1 wherein tumor cells previously isolated from said patient provide the tumor antigens.
3. The method of claim 1 wherein the cytokine gene is selected from the group consisting of interleukin-1, interleukin-2, interleukin-3, interleukin-4, interleukin-5, interleukin-6, and gamma-interferon.
4. The method of claim 3 wherein one cytokine gene is interleukin-2.
5. The method of claim 1 wherein at least one cytokine gene is transferred into cells to generate CE cells by recombinant methods.
6. The method of claim 5 wherein the cytokine gene is present in an expression vector.
7. The method of claim 6 wherein said expression vector additionally contains a suicide gene.
8. The method of claim 5 wherein the CE cells are generated from fibroblasts and antigen-presenting cells.

9. A method for enhancing a patient's immune response to a cancer comprising:

- a) isolating fibroblasts from said patient;
- b) culturing said fibroblasts in vitro;
- c) transducing said fibroblasts with a retroviral expression vector containing the gene coding for IL-2 and a gene coding for a tumor antigen in a retroviral expression vector, to express said tumor antigen and to express and secrete said IL-2 by said fibroblasts; and
- d) immunizing said patient with said fibroblasts that express IL-2 at a level sufficient to enhance an immune response but low enough to avoid substantial systemic toxicity and that express said tumor antigen, at a site other than an active tumor site.

10. The method of claim 9 wherein said fibroblasts are further modified to express a suicide gene.

11. A composition for increasing a patient's immune response to tumor antigens comprising tumor antigens and CE cells genetically modified to express at least one cytokine gene product.


12. The composition of claim 11 wherein the cytokine gene is selected from the group consisting of interleukin-1, interleukin-2, interleukin-3, interleukin-4, interleukin-5, interleukin-6, and gamma interferon.

13. The composition of claim 12 wherein one cytokine gene is interleukin-2.


14. The composition of claim 11 wherein each cytokine gene is expressed at a level sufficient to stimulate the immune response but low enough to avoid substantial systemic toxicities.

15. The method of claim 9 wherein in said transducing step said retroviral expression vector has a promoter causing sustained secretion of IL-2.


16. The method of claim 15 wherein said retroviral expression vector causes the secretion of at least four units of IL-2 per day for a period of ten days or longer.



RETROVIRAL VECTOR DC/TKIL2



RETROVIRAL VECTOR LXSN-IL2



RETROVIRAL VECTOR LNCX-IL2

FIG. 1

2 / 15




FIG. 2

3 / 15




FIG. 3

4 / 15





FIG. 4



1 2 3 4 5 6 7

FIG. 5

6 / 15




FIG. 6

7 / 15




FIG. 7

8 / 15




FIG. 8

9 / 15




FIG. 9

10 / 15




FIG. 10

SUBSTITUTE SHEET




FIG. 11

12 / 15




FIG. 12

SUBSTITUTE SHEET




FIG. 13

14 / 15




FIG. 14

SUBSTITUTE SHEET

15 / 15




FIG. 15

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US92/08999

A. CLASSIFICATION OF SUBJECT MATTER

IPC(S) : Please See Extra Sheet.

US CL : Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/93B, 93U, 89; 435/240.2, 320.1, 69.5, 69.51, 69.52; 935/65, 32, 12, 57, 70, 71

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

BIOSIS, MEDLINE, APS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X Y	Journal of Experimental Medicine, Volume 172, issued October 1990, Gansbacher et al., "Interleukin 2 Gene Transfer into Tumor Cells Abrogates Tumorigenicity and Induces Protective Immunity", pages 1217-1224, see the entire document.	<u>1-8, 11-14</u> 9, 10, 15, 16
X Y	Cell, Volume 57, issued 05 May 1989, Tepper et al., "Murine Interleukin-4 Displays Potent Anti-Tumor Activity In Vivo", pages 503-512, see the entire document.	<u>1-3, 5, 6, 8, 11, 12, 14</u> 4, 13
X Y	Cell, Volume 60, issued 09 February 1990, Fearon et al., "Interleukin-2 Production by Tumor Cells Bypasses T Helper Function in the Generation of an Antitumor Response", pages 397-403, see the entire document.	<u>1-3, 5, 8, 11-13</u> 2, 6, 7, 14-16
Y	Cancer Research, Volume 50, issued 15 August 1990, Ogura et al., "Implantation of Genetically Manipulated Fibroblasts into Mice as Antitumor α -Interferon Therapy", pages 5102-5106, see the entire document.	1-16

☒ Further documents are listed in the continuation of Box C.
 ☐ See patent family annex.

* "A"	Special categories of cited documents: document defining the general state of the art which is not considered to be part of particular relevance	"T"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"E"	earlier document published on or after the international filing date	"X"	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L"	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y"	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O"	document referring to an oral disclosure, use, exhibition or other means		
"P"	document published prior to the international filing date but later than the priority date claimed	"Z"	document member of the same patent family

Date of the actual completion of the international search

11 January 1993

Date of mailing of the international search report

26 JAN 1993

 Name and mailing address of the ISA/
 Commissioner of Patents and Trademarks
 Box PCT
 Washington, D.C. 20231

Authorized officer

JACQUELINE STONE

Facsimile No. NOT APPLICABLE

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US92/08999

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X Y	Cancer Research, Volume 50, issued 15 December 1990, Gansbacher et al., "Retroviral Vector-mediated Interferon Gene Transfer into Tumor Cells Generates Potent and Long Lasting Antitumor Immunity", pages 7820-7825, see the entire document.	1, 3, 5, 6, 8, 11, 12, 14 2, 7

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US92/08999

A. CLASSIFICATION OF SUBJECT MATTER:

IPC (5):

A61K 48/00, 35/12, 39/00; C12N 15/19, 15/24, 15/25, 15/26, 15/90, 15/63

A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

424/93B, 93U, 89; 435/240.2, 320.1, 69.5, 69.51, 69.52; 935/65, 32, 12, 57, 70, 71